

RECORDS
of the
INDIAN MUSEUM

Vol. XXV, Part IV, pp. 399—403.

Note on the significant characters of local variation
in proportion of dextral and sinistral shells
in samples of the snail *Bulliminus*
dextrosinister.

By
P. C. MAHALANOBIS.

CALCUTTA:
AUGUST, 1923.

STATISTICAL NOTE ON THE SIGNIFICANT CHARACTER OF
LOCAL VARIATION IN PROPORTION OF DEXTRAL AND
SINISTRAL SHELLS IN SAMPLES OF THE SNAIL *BULIMINUS*
DEXTROSINISTER FROM THE SALT RANGE, PUNJAB.

By P. C. MAHALANOBIS, B.Sc., M.A. (Cantab.), Professor of Physics,
Presidency College, Calcutta.

Dr. Annandale, Director, Zoological Survey of India, sent me the following data for a statistical report on a collection of shells of the snail *Buliminus dextrosinister* from the Salt Range.

Locality.	DEXTRAL.		SINISTRAL.		TOTAL
	No.	Percentage.	No.	Percentage.	
1. Chalisa	2	100	2
2. Katas	10	83.3	2	16.7	12
3. Kallar Kahar	21	33.3	49	66.7	70
4. Sardhi	2	4.2	46	95.8	48
TOTAL	35	37.7	97	62.3	132

Two distinct questions arise.

- (a) Do dextral and sinistral shells occur in nature in equal proportions or do they occur in different proportions ?
- (b) If in different proportion, is this proportion the same for all localities, or is there local variation in the proportion ?

Let us assume that the shells occur in equal proportions in nature. Then the "theoretical proportion" of each is $\frac{1}{2}$. For example in the total sample, the "theoretical proportion" would be 66 of each. The "observed proportion" is 35 dextral and 97 sinistral.

Of course we cannot expect to get 66 dextral shells in each sample of 132. The "observed proportions" will vary on account of fluctuations of sampling. The question is whether difference between the "observed" and the "theoretical" proportion is merely due to such fluctuations of sampling or whether it is indicative of a real difference in the proportions in which dextral and sinistral shells occur in nature.

To put it in a slightly different way : A sample of 35 dextral and 97 sinistral shells may sometimes occur on account of fluctuations of sampling. But how often ? In other words, the precise mathematical question is : What is the exact probability of such occurrence ?

The solution is well-known and may be stated quite generally.

If the "theoretical proportion" of dextral and sinistral shells is p and q respectively ($p + q = 1$) then the probability of occurrence of a sample of s dextral and r sinistral ($s + r = m$) in a sample of m is given by

$$C_s = \frac{m!}{s! r!} \cdot (p)^s (q)^r$$

By direct arithmetical calculation I obtain the following table.

TABLE II.—Probability of occurrence of "observed" samples for $p=q=\frac{1}{2}$.

Locality.	Number of dextral shells.	Probability.	odds against
1. Chalisa	2	0.25	4 to 1
2. Katas	10 or more.	0.01 92 86	51 to 1
3. Kallar Kahar	21 or less	5.46 $\times 10^{-4}$	18 28 to 1
4. Sardhi	2 or less	4.16 21 $\times 10^{-112}$	24.10 ¹⁸ to 1
TOTAL	35 or less	3.15 27 $\times 10^{-9}$	32.10 ⁶ to 1

In the case of *Chalisa* the evidence is quite inconclusive. *Katas* also is doubtful; odds of 51 to 1 are not sufficiently high to justify us in asserting that the proportion of dextral shells is really greater than $\frac{1}{2}$. On the other hand the odds are very considerable in the case of *Kallar Kahar* and practically overwhelming in the case of the *Sardhi* and the *Total*. Thus on the whole we may reasonably argue that dextral shells occur in substantially lower proportion in nature.

I may now pass on to the second question, namely, is there any local variation in the proportion? The problem may be stated quite generally as follows.

In a first sample of n shells, p are found to be dextral and q sinistral ($p + q = n$), what is the chance of occurrence of s dextral and r sinistral shells in another sample of m ($m = r + s$)?

The solution is given by Bayes' theorem and the result can be easily calculated with the help of formulae given by M. Greenwood¹ and Karl Pearson.²

$$C_s = \frac{\binom{n+1}{p} \binom{m}{s}}{\binom{p}{p} \binom{q}{q} \binom{m+n+1}{m}} \cdot \frac{\binom{p+s}{s} \binom{q+r}{r}}{\binom{p}{p} \binom{q}{q} \binom{m}{m}}$$

Greenwood³ says "Let the chance of a_2 or more successes in n_2 (s or more successes in m_1 in our notation) after a_1 successes in n_1 (i.e. p successes in n) be p_2 and the chance of a_1 or less successes in n_1 trials (p or less in n) after a_2 successes in n_2 (s success in m) be p_2 . Then, since either n_1 or n_2 might have been drawn first, a measure of the probability of the observed result will be $\frac{1}{2} (p_1 + p_2)$."

¹ *Biometrika*, IX, pp. 69-90.

² *Phil. Mag.* 1907, p. 365 also *Biometrika*, XIII, 1920, pp. 1-16, and *Tables for Statisticians and Biometricians*, pp. Ixx-Ixxiii.

³ *Ibid.* p. 81.

I am not quite sure about the validity of Greenwood's argument, at least, in its application to the present problem. It seems desirable to give full weight to the most favourable probability for agreement before differentiation is definitely asserted. On this principle the more numerous sample should invariably be taken as the first sample. Doing this we shall find the most favourable probability that the samples are in agreement. If this most favourable probability itself is found to be too small then we shall be justified in asserting differentiation. In other words we should choose the case most unfavourable for our conclusion, thus erring on the safe side.

I have also used the "four-fold X^2 method" of Pearson¹ as a check. In the present notation.

$$X^2 = \frac{(pr - qs) \cdot 2 (m+n)}{n \cdot m \cdot (p+s) (q+r)}$$

and may be easily calculated. The probability of occurrence, P, is then found from Tables XII and XVII (pp. 26, 31) of Tables for Biometricians and Statisticians. Q gives the probability on Bayes' theorem.

Chalisa : Katas

Q = .6286	6 to 10 in favour of agreement.
P = .312	1 to 3 in favour of agreement.

There is no evidence of differentiation. The probability is that both represent the same content in nature.

Chalisa : Kallar Kahar

Q = .0927	94 to 10 against agreement.
P = .2271	44 to 10 against.

Differentiation is not improbable but the evidence is not conclusive.

Chalisa : Sardhi

Q = .0047	212 to 1 against.
P = 2.55×10^{-5}	40,000 to 1 against.

Differentiation is probable; but the smallness of the number (2 only) in the case of Chalisa renders the conclusion somewhat doubtful.

Katas : Kallar Kahar

Q = .00065	1528 to 1 against.
P = .00630	157 to 1 against.

It seems fairly certain that Katas has got a higher proportion of dextral shells than Kallar Kahar.

Kallar Kahar and Sardhi

Q = .000177	5662 to 1 against.
P = .007083	140 to 1 against

These also seem to be quite significantly differentiated.

¹ *Draper's Company Research Memoirs, Biometric Series VIII.* "On a Novel Method of regarding the Association of two varieties classed solely in Alternate Categories."

Katas : Sardhi

Q = 5.5.10⁻⁶ 180,800 to 1 against.

P = 1.6.10⁻⁹ 10⁶ to 1 against.

The probability is now overwhelming that the samples are different.

Summing up we may say : Chalisa is uncertain ; Katas shows distinctly higher proportion of dextral shells than either Kallar Kahar or Sardhi, while Kallar Kahar itself shows significant excess over Sardhi.

The following table will give some idea about the reliability of the observed proportion in each case :—

TABLE III.—Percentage frequency of dextral shells in samples of the same size *m*.

1. Chalisa <i>m</i> = 2	3. Kallar Kahar <i>m</i> = 70	4. Sardhi <i>m</i> = 48	Total <i>m</i> = 132
0 10.00	10 0.679	0 13.350	22 0.9137
1 30.00	11 1.087	1 19.143	23 1.2244
2 60.00	12 1.634	2 19.349	24 1.5927
	13 2.316		25 2.0137
	14 3.113		26 2.4781
	15 3.985	3 16.124	27 2.9713
	16 4.874	4 11.960	28 3.4571
	17 5.711	5 8.186	29 3.9682
	18 6.430	6 5.273	30 4.4278
	19 6.970	7 3.236	31 4.8319
	20 7.287	8 1.962	32 5.1606
	21 7.360	9 1.084	33 5.3979
		10 0.597	34 5.5332
			35 5.5616
	22 7.193		36 5.4843
	23 6.809		37 5.3085
	24 6.251		38 5.0457
	25 5.569		39 4.7119
	26 4.819		40 4.3244
	27 4.054		41 3.9019
	28 3.316		42 3.4626
	29 2.638		43 3.0230
	30 2.043		44 2.5971
	31 1.540		45 2.1962
	32 1.131		46 1.8285
	33 0.808		47 1.4992
			48 1.2107
			49 0.9632

Taking 1% (one per cent of the samples) as a limiting value, we find the range to be 0 to 2 (i.e. 0% to 100%) dextral shells in the case of Chalisa, 5 to 12 (i.e. 42% to 100%) in the case of Katas, 11 to 32 (16% to 46%) in the case of Kallar Kahar, 0 to 9 (0% to 19%) in the case of Sardhi and 23 to 48 (17% to 29%) in the case of the total sample.

Finally we may compare each sample with the three others taken together.

		Dextral.	Sinistral.	Total.
1. Others	...	33	97	130
Chalisa	...	2	0	2
		35	97	132

P = .0678. The evidence is inconclusive.

2. *Katas and others*

Others	...	25	95	120
Katas	...	10	2	12

$P=2.38 \times 10^{-5}$ Katas seems to be really differentiated from the rest.

3. *Kallar Kahar and others*

Kallar Kahar	...	21	49	70
Others	...	14	48	62

$P=.1863$ Kallar Kahar agrees well with the rest of the sample and may be considered fairly typical.

4. *Sardhi and others*

Sardhi	...	2	46	48
Others	...	33	51	84

$P=2.38 \times 10^{-5}$ Sardhi also is different from the rest.

Conclusion.—We may on the whole conclude that dextral shells occur in less frequency in nature, the proportion probably being roughly from a fifth to a third; that Kallar Kahar is a fairly typical sample, while Katas and Sardhi seem to be somewhat differentiated from the rest.