

# Ratio-dependent predator–prey model: effect of environmental fluctuation and stability

M Bandyopadhyay<sup>1</sup> and J Chattopadhyay<sup>2,3</sup>

<sup>1</sup> Department of Mathematics, Scottish Church College, Kolkata 700006, India

<sup>2</sup> Embryology Research Unit, Indian Statistical Institute, Kolkata 700108, India

E-mail: malay\_ban@rediffmail.com and joydev@isical.ac.in

Received 27 April 2004, in final form 11 November 2004

Published 1 February 2005

Online at [stacks.iop.org/Non/18/913](http://stacks.iop.org/Non/18/913)

Recommended by J A Glazier

## Abstract

This paper deals with the problem of a ratio-dependent predator–prey model. The deterministic and stochastic behaviour of the model system around biologically feasible equilibria are studied. Conditions for which the deterministic model enter into Hopf-bifurcation are worked out. Stochastic stability of the system around positive interior equilibrium is studied. To substantiate our analytical findings numerical simulations are carried out for a hypothetical set of parameter values.

Mathematics Subject Classification: 34D05, 34K20, 92D25

## 1. Introduction

The dynamical relationship between prey and their predators has long been and will continue to be one of the dominant themes in ecology due to its universal existence and importance [16, 18]. The dynamical problems involved with mathematical modelling of predator–prey systems may appear to be simple at first sight; however, the detailed analysis of these model systems often leads to very complicated as well as challenging problems. The most important part of modelling the population ecosystem is to make sure that the concerned mathematical model can exhibit well-known system behaviour for the system under consideration. Dynamical modelling of ecological systems is a frequently evolving process. A systematic mathematical approach can lead to a better understanding of the plausible models and the exposed discrepancies in turn lead to the necessary modifications [43].

After the pioneering work of Alfred Lotka and Vito Volterra in the middle of 1920 for predator–prey interactions, prey-dependent predator–prey models were studied extensively

[26, 31, 49, 50]. In population dynamics, a functional response of the predator to prey density refers to the change in the density of prey per unit time per predator as a function of the prey density. Quite a good number of works have already been performed in ecological systems [7, 10, 14, 21, 23, 33, 62] where the model systems are based on prey-dependent model systems. The classical prey–predator models with prey-dependent functional response take the form (or some equivalent form)

$$\frac{dN}{dt} = rN \left(1 - \frac{N}{K}\right) - p(N)P, \quad \frac{dP}{dt} = cp(N)P - eP, \quad (1.1)$$

where  $p(N)$  is the so-called prey-dependent functional response ( $N(t)$  and  $P(t)$  denote prey and predator population density at any instant of time ‘ $t$ ’, respectively, and other parameters have the usual meaning). In most cases prey-dependent functional response  $p(N)$  is given by  $p(N) = aN/(b + N)$  or  $p(N) = aN^2/(b + N^2)$  or  $p(N) = aN^2/(b + N + \alpha N^2)$  or  $p(N) = aN^\theta/(b + N^\theta)$  or some equivalent form. These types of classical predator–prey model systems exhibit not only the wellknown ‘paradox of enrichment’ formulated by Hairston *et al* [34] and Rosenzweig [56] but also the so-called ‘biological control paradox’, which was introduced by Luck [47]. The ‘paradox of enrichment’ states that enriching a predator–prey system (by increasing the carrying capacity) will cause an increment in equilibrium density of predator but not of prey population which in turn destabilizes the interior equilibrium. As a result it increases the chance of stochastic extinction of the predator population. However, in nature, it is observed that enriching the system increases the prey density which is not a factor responsible for destabilizing a stable equilibrium and fails to increase the amplitude of oscillations in systems that already cycles [1].

The so-called ‘biological control paradox’ states that we cannot have a low and stable prey equilibrium density, which is in contradiction with many examples of successful biological controls where the prey population is maintained at low densities compared to its environmental carrying capacity [4, 6]. A further example is, cactoblastis–opuntia in Australia, where the crucial factor seems to be pseudo-interference (see, May [49]), and where biological control has worked and resulted in low and stable pest densities. So this paradox is a pure artefact, created by simplifying assumptions on functional response. For the rest, the paradox of enrichment exists, but only in systems where one predator–prey pair exists in isolation and the predictions radically change when they are embedded in a simple food chain model [52]. Most of the natural systems are indeed very complex with the predators being exposed to various degrees of facultative secondary carnivory and interacting with many other subsystems in various ways. This indicates the fact that the paradox of biological control is not intrinsic for most predator–prey systems.

There are some good data on simple as well as fairly isolated predator–prey systems, like Mary Power’s catfish–alga system in the Panamanian streams [54] where one can find the interesting phenomenon that some increment in the resource for prey population at equilibrium increases the predator density in place of prey abundance. In some predator–prey systems where the Holling disc is a reasonable assumption for predators (e.g. the weasel–vole system found in boreal Europe, where voles have a hiding place and weasels have no alternative resources), we observe large amplitude oscillations of population distribution like carnivore–herbivore systems [52, 53]. According to the ‘paradox of enrichment’ these types of large amplitude oscillations are expected in relatively productive areas and in the most reproductive parts of tundra, while voles inhabiting less productive tundra areas are relatively stable [55]. High amplitude oscillations were the basic characteristics in high alpine barrens where the density of predator is low, but at present, rodent time trajectories indicate the fingerprints of predators [60]. The change in dynamic position of herbivores along productivity

gradients thus indicates the necessity of reasonable alteration or corrections of simple as well as traditional food chain models [52]. Recently, predator–prey models with prey-dependent response function have been facing a great challenge from biological and physiological researchers [3–5, 32]. At present it is clear that predator abundance also has the ability to influence the functional response. Arditi and Ginzburg [5] have suggested that, in situations characterized by strong space and time heterogeneities, the functional response can be approximated by a function of the prey-to-predator ratio. Several biologists have been able to establish the fact that functional responses over typical ecological timescales ought to depend on the densities of both prey and predators, especially when predators have to search for food and therefore have to share or compete for food. Actually prey-dependent and ratio-dependent models are extremes or limiting cases; prey-dependent models are based on the daily energy balance of predators, on the other hand ratio-dependent models presuppose that prey are easy to find and that predator dynamics are, in essence, governed by direct density dependence, with prey densities determining the sizes of defended territories. Within the natural environment both aspects have the ability to influence predator–prey dynamics, and the issue of which of the two extremes is closer to reality in which system is wide open. Moreover, the pursuit of the prey-dependent approach has proved more fertile since its ‘paradoxes’ seem to be quite realistic where the premises for their existence are found, but here opinions may differ and there is no strong evidence to close one door or another.

Arditi and Ginzburg [5] first proposed the following Michaelis–Menten–Holling type, ratio-dependent predator–prey system

$$\frac{dN}{dt} = rN \left(1 - \frac{N}{K}\right) - \frac{\alpha NP}{k_1 P + N}, \quad \frac{dP}{dt} = \frac{c\alpha NP}{k_1 P + N} - eP. \quad (1.2)$$

The dynamics of this type of ratio-dependent predator–prey model has been studied by many researchers (see, e.g., Kuang and Beretta [44], Jost *et al* [39], Hsu *et al* [38]).

Major parts of the work in this direction are based on deterministic models of differential and difference equations. The deterministic approach has, however, some limitations in biology: it is always difficult to predict the future of the system accurately. Deterministic models in ecology do not usually incorporate environmental fluctuations based on the idea that in the case of large populations, stochastic deviations (or the effect of random environmental fluctuations) are small enough to be ignored. A stochastic model provides a more realistic picture of a natural system than its deterministic counterpart. A central obstacle in the stochastic modelling of an ecosystem is the lack of mathematical machinery available to analyse nonlinear multi-dimensional stochastic models [28, 40, 41]. A quantum leap in the mathematical sophistication of ecological modelling occurred when May [49] introduced stochastic differential equations (SDEs) to investigate limits to niche overlap in a randomly fluctuating environment. Well-known deterministic population models (such as the Lotka–Volterra model, Gauss type prey–predator model, etc) are the starting points of stochastic multi-species models which include demographic or environmental stochasticity. The resulting stochastic models involve nonlinear SDEs whose solutions pose great difficulties. Different techniques of linearization of nonlinear SDEs giving rise to a set of deterministic moment equations have been receiving a great deal of attention in different fields of science and technology [8, 11, 12, 28, 29, 40, 41, 51, 57].

The main objective of this paper is to consider a ratio-dependent predator–prey model and its stability behaviour around different equilibrium points with special emphasis on the controversial equilibrium point (0, 0) for the ratio-dependent model. Our next objective is to develop a stochastic dynamic model for the ratio-dependent predator–prey model and to examine the stability of the model system under random environmental fluctuations. We make

a comparative analysis of the stability of the model system within deterministic and stochastic environments.

## 2. Deterministic model

The classical model for predator–prey systems can be written in its classical form by a system of first order nonlinear ordinary differential equations as

$$\frac{dN}{dt} = Nf(N) - g(N, P)P, \quad \frac{dP}{dt} = h(N, P)P - \gamma P \quad (2.1)$$

with prey abundance  $N(t)$  and predator abundance  $P(t)$  at any given instant of time ‘ $t$ ’.  $f(N)$  is the per capita rate of increase of the prey in the absence of predation and ‘ $\gamma$ ’ is the food-independent mortality rate of the predator, assumed to be constant. The amount of prey biomass consumed by each predator per unit of time is described by the function  $g(N, P)$ , while  $h(N, P)$  describes per capita production rate of the predator. There is a considerable amount of evidence to show that predator production rate can be modelled as simply proportional to food intake (up to a very good approximation)

$$h(N, P) = e g(N, P), \quad (2.2)$$

where the constant ‘ $e$ ’ is interpreted as conversion efficiency and satisfies the condition  $0 < e < 1$ . Specific examples are given by Slobodkin [58] for hydras, by Beddington *et al* [13] for numerous arthropods and by Coe *et al* [24] for large African herbivores in support of previous assumptions. The trophic function  $g(N, P)$  is the sole link between prey–predator dynamics [5]. In this paper we will consider the usual logistic form of the growth function for prey in the absence of predator as

$$f(N) = r \left( 1 - \frac{N}{K} \right),$$

where ‘ $r$ ’ is the intrinsic growth rate of prey and ‘ $K$ ’ is the environmental carrying capacity. For the traditional prey-dependent predator–prey models (e.g. (1.1)), the functional response  $g(N, P)$  depends only on the density of prey population ‘ $N$ ’ as we have remarked in the introduction of this paper. According to Berryman [18], credible and consistent predator–prey models must be able to satisfy the minimum biological property that ‘when resources are low relative to population density, the predator per-capita growth rate should decline with its density’. However, the prey-dependent predator–prey models are unable to satisfy this criterion due to the fact that the predator per-capita growth rate becomes a function of prey population ‘ $N$ ’ only, which is independent of the density of predators ‘ $P$ ’. To overcome this situation, Arditi and Ginzburg [5] have suggested that for situations characterized by strong space and time heterogeneities, the functional response can be approximated by a function of the prey-( $N$ )-to-predator-( $P$ ) ratio ( $N/P$ ) and this leads to the ratio-dependent functional response  $g(N, P)$  as follows:

$$g := g \left( \frac{N}{P} \right) = \frac{\alpha(N/P)}{1 + \alpha\beta(N/P)} = \frac{\alpha N}{P + \alpha\beta N},$$

where ‘ $\alpha$ ’ represents the total attack-rate for predator and ‘ $\beta$ ’ is the handling time and the functional response is known as the Michaelis–Menten–Holling type functional response [17, 39, 43]. Under these assumptions the dynamics of the ratio-dependent predator–prey system is governed by the following system of first order nonlinear ordinary differential equations:

$$\frac{dN}{dt} = rN \left( 1 - \frac{N}{K} \right) - \frac{\alpha NP}{P + \alpha\beta N}, \quad \frac{dP}{dt} = \frac{e\alpha NP}{P + \alpha\beta N} - \gamma P. \quad (2.3)$$

Initial conditions for the system of equation (2.2) are given by  $N(t = 0) = N_0 \geq 0$  and  $P(t = 0) = P_0 \geq 0$ . It is typical for predator–prey systems: positive  $N$ -axis, positive  $P$ -axis and the interior of the first quadrant invariant under system (2.3) and the solutions with positive initial condition continue to be positive for all time ‘ $t$ ’. As observed by Freedman and Mathsen [27], Jost *et al* [39], Kuang and Beretta [44] and Kuang [43], system (2.3) is not well-defined at the origin  $(0, 0)$  and hence the model is unable to capture the idea that the growth rate of both the populations are zero in the absence of prey as well as predator population. To overcome this situation, Xiao and Ruan [63] have modified the model system (2.3) by redefining it as follows:

$$\frac{dN}{dt} = rN \left( 1 - \frac{N}{K} \right) - \frac{\alpha NP}{P + \alpha\beta N}, \quad \frac{dP}{dt} = \frac{e\alpha NP}{P + \alpha\beta N} - \gamma P \quad \text{when } (N, P) \neq (0, 0), \tag{2.4a}$$

$$\frac{dN}{dt} = 0 = \frac{dP}{dt} \quad \text{when } (N, P) = (0, 0). \tag{2.4b}$$

For the sake of simplicity it is convenient to scale the variables as  $x = N/K, y = P/K\alpha\beta$  and consider the dimensionless time  $\tau = tr$ . The dimensionless equations are then given by

$$\frac{dx}{d\tau} = x(1 - x) - \frac{axy}{x + y} \equiv F_1(x, y), \quad \frac{dy}{d\tau} = \frac{bxy}{x + y} - cy \equiv F_2(x, y), \tag{2.5a}$$

$$F_1(0, 0) = F_2(0, 0) = 0, \tag{2.5b}$$

where  $a = \alpha/r, b = e/\beta r$  and  $c = \gamma/r$ . For convenience, in the following, time  $\tau$  is replaced by  $t$  as the dimensionless time. The initial conditions for the system of equations (2.5) are given by  $x(0) = x_0 \geq 0$  and  $y(0) = y_0 \geq 0$  which is also biologically meaningful.

2.1. Boundedness

Due to the boundedness of the functional responses, we see that

$$\lim_{(x,y) \rightarrow (0,0)} F_1(x, y) = \lim_{(x,y) \rightarrow (0,0)} F_2(x, y) = 0.$$

Using equation (2.5b) we can conclude that the functions  $F_1(x, y)$  and  $F_2(x, y)$  are continuous functions on  $\mathbb{R}_+^2 = [(x, y) : x \geq 0, y \geq 0]$ . Straightforward computation shows that they are Lipschitzian on  $\mathbb{R}_+^2$ . Hence a solution of (2.5) with non-negative initial condition exists and is unique. It is also easy to see that these solutions exist for all  $t > 0$  and stay non-negative. In fact, if  $x(0) = x_0 > 0$ , then  $x(t) > 0$  for all  $t > 0$ . The same argument is true for the  $y$ -component. Hence, the interior of  $\mathbb{R}_+^2$  is invariant under model system (2.5). Our next task is to consider the boundedness for the solutions of the model system (2.5).

**Lemma 1.** *All the solutions of the system (2.5) with the positive initial condition  $(x_0, y_0)$  are uniformly bounded within a region  $\Omega$ , where,*

$$\Omega = \left\{ (x, y) : 0 \leq x \leq 1, 0 \leq x + \frac{a}{b}y \leq \frac{L}{c} \right\}$$

with  $L = c + 1/4$ .

**Proof.** From the first equation of (2.5) we get,

$$\frac{dx}{dt} = x(1 - x) - \frac{axy}{x + y} \leq x(1 - x).$$

So,  $x(t) \leq 1$ , as  $t \rightarrow +\infty$ .

Let us define the function

$$W(t) = x(t) + \frac{a}{b}y(t).$$

Calculating the time derivative of  $W(t)$  along the trajectories of equations (2.5), we get

$$\frac{d}{dt}W(t) = \frac{dx}{dt} + \frac{a}{b}\frac{dy}{dt} = x(1-x) - \frac{ac}{b}y.$$

Clearly, the maximum value of  $x(1-x)$  is  $1/4$ , whenever  $0 \leq x \leq 1$ . Then,

$$\frac{d}{dt}W(t) + cW(t) = x(1-x) + cx \leq c + \frac{1}{4} = L \quad (\text{say}).$$

Thus, as  $t \rightarrow +\infty$ ,  $0 \leq W(t) \leq L/c$  [19]. Hence system (2.5) is dissipative with the asymptotic bound  $L/c$ . This ensures the existence of a compact neighbourhood  $\Omega$  which is a proper subset of  $\mathbf{R}_+^2$  such that for sufficiently large initial conditions  $(x_0, y_0)$  the trajectories of the system of equations (2.5) will always be within the set  $\Omega$ .

Hence we have shown that the model system (2.5) is dissipative.

## 2.2. Equilibria

For population models in deterministic environments, with the environmental parameters being well-defined constants, it is natural curiosity to find the community equilibria where all the species' populations have time independent values, that is, where all net growth rates are zero. Classical two species predator-prey models always possess at least three equilibrium points: (i) trivial equilibrium, (ii) axial equilibrium and (iii) positive interior equilibrium [45]. Earlier works on ratio-dependent predator-prey models (e.g. [27, 39, 43, 44]) have mentioned that the model system (2.3) cannot be linearized at  $(0, 0)$  and consistent dynamical analysis for the model system in the vicinity of the origin reveals the rich and complicated dynamics. Based on the redefined ratio-dependent predator-prey model (2.4) and its non-dimensionalized version (2.5) we are able to mention that the system of equations (2.5) has two equilibria,  $E_0(0, 0)$  and  $E_1(1, 0)$  on the  $x$ -axis for all possible as well as admissible values of the parameters involved with the model system (2.5). The third and most interesting equilibrium point (from a biological point of view) is  $E_+(x^*, y^*)$  where  $x^*$  and  $y^*$  are non-zero positive solutions of the equations  $F_1(x, y) = 0 = F_2(x, y)$  and are given by

$$x^* = 1 - \left[ \frac{a(b-c)}{b} \right] \quad \text{and} \quad y^* = \left[ \frac{(b-c)x^*}{c} \right]. \quad (2.6)$$

Simple mathematical argument shows that  $x^*$  is positive for all  $a < b/(b-c)$  and positivity of  $y^*$  demands an extra condition  $b > c$ .

## 2.3. Behaviour around $E_0(0, 0)$

The study of stability plays a significant role in understanding the structure and functions of ecological systems. A variety of ecologically interesting interpretations are involved with the term 'stability'. The most common meaning corresponds to *neighbourhood stability* which means stability in the vicinity of an equilibrium point associated with the deterministic model system. An equilibrium point is called a stable equilibrium point if, when the populations are perturbed they will return to the equilibrium point with the advancement of time. The return may be achieved either as damped oscillations or monotonically. It is common practice to find the linearized system of equations around the equilibrium point and the sign of eigenvalues associated with the corresponding Jacobian matrix determines the stability around the equilibrium point. Before studying the stability of controversial equilibrium point  $E_0(0, 0)$

we would like to remark that mathematical investigation of the nature of the solution for the model system in the vicinity of origin reveals some interesting dynamical behaviour depending on the parametric restrictions. This type of dynamical behaviour has never been observed in classical prey-dependent predator–prey models. Based on the outcome of stability analysis around  $E_0(0, 0)$ , one can obtain a clear idea of the possibility of extinction of population at low population density of both prey and predators. In the case of the ratio-dependent predator–prey model it is not possible to linearize the model system around  $E_0(0, 0)$ . To study the behaviour of the model system around  $E_0$  we follow the technique developed by Arino *et al* [7]. (For rigorous calculations and mathematical justification of the said technique, interested readers may consult the work of Arino *et al* [7] and the references therein.) For this purpose the model system (2.5) can be written as

$$\frac{dX}{dt} = H(X(t)) + Q(X(t)), \tag{2.7}$$

where  $H(\cdot)$  is  $C^1$  except at the origin, is a continuous and homogeneous function of degree one, i.e.  $H(sX) = sH(X)$  for all scalar  $s \geq 0$ ,  $X(t) = (x_1, x_2) \in \mathbb{R}_+^2$  and  $Q$  is a  $C^1$  function and satisfies the condition  $Q(X) = o(X)$  in the vicinity of the origin. The functions  $H(\cdot)$  and  $Q(\cdot)$  are defined by

$$H(X) \equiv (H_1(X), H_2(X)), \quad Q(X) \equiv (Q_1(X), Q_2(X)), \tag{2.8}$$

$$H_1(X) = x_1 - \frac{ax_1x_2}{x_1 + x_2}, \quad H_2(X) = \frac{bx_1x_2}{x_1 + x_2} - cx_2, \tag{2.9}$$

$$Q_1(X) = -x_1^2, \quad Q_2(X) = 0. \tag{2.10}$$

Let us assume that  $X(t)$  is a solution of (2.7), is bounded such that  $\liminf_{t \rightarrow \infty} \|X(t)\| = 0$ . For this situation it is possible to find a sequence  $X(t_n + \cdot) \rightarrow 0$  uniformly as  $t_n \rightarrow \infty$ . Define

$$y_n(s) = \frac{X(t_n + s)}{\|X(t_n + s)\|}. \tag{2.11}$$

Clearly,  $Y_n$  is a sequence with  $\|y_n\| = 1$ . Applying the Ascoli–Arzela theorem [20], it is possible to find a subsequence corresponding to  $y_n$  which converges towards some function  $y(t)$  and the limiting function satisfies the equation

$$\frac{dy}{dt} = H(y(t)) - (y(t), H(y(t)))y(t). \tag{2.12}$$

In equation (2.12),  $(\cdot, \cdot)$  stands for the standard inner-product in  $\mathbb{R}^2$ . The steady states of (2.12) are vectors  $V$  satisfying  $H(V) = (V, H(V))V$  which are solutions of the nonlinear eigenvalue problem,  $H(V) = \mu V$  with  $\mu = (V, H(V))$ . These stationary solutions correspond to the fixed directions along which the trajectories of (2.12) may converge asymptotically. Using (2.9), the nonlinear eigenvalue problem becomes

$$[(1 - \mu)v_1 + (1 - a)v_2]v_1, \quad [(b - c)v_1 - (c + \mu)v_2]v_2 = 0. \tag{2.13}$$

Now we are in a position to discuss in detail the possibility of reaching the origin following fixed directions.

*Case 1.*  $v_1 = 0$  and  $v_2 \neq 0$

In this case, there is a possibility of reaching the origin along the  $y$ -axis, with  $\mu = -c$ .

*Case 2.*  $v_1 \neq 0$  and  $v_2 = 0$

In this case, it is possible to reach the origin along the  $x$ -axis, with  $\mu = 1$ .

Case 3.  $v_1 \neq 0$  and  $v_2 \neq 0$

In this case, the possibility of reaching the origin along some fixed direction from the interior of the first quadrant depends on the existence of a real root of the quadratic equation

$$\mu^2 + \mu(c - 1) + ab - b - bc = 0. \quad (2.14)$$

The existence of a real root of the quadratic equation (2.14) demands

$$c < b < \frac{1}{a-1} \left[ \frac{(c-1)^2}{4} + ac \right]. \quad (2.15)$$

Again the upper bound for the parameter  $b$  will be biologically feasible if and only if  $a > 1$  or equivalently  $\alpha > r$ . This result has the biological significance that if the prey-catching capacity is higher than the intrinsic growth rate of predator then both the populations approach total extinction.

Under the three conditions discussed above, it is possible to reach the trivial equilibrium point  $E_0(0, 0)$  and hence  $E_0$  is an attractor for the model system (2.5).

#### 2.4. Behaviour around $E_1$ and $E_*$

The Jacobian matrix  $J(x, y)$  for the system (2.5) at any point of the first quadrant  $(x, y)$ , except at the origin, is given by

$$J(x, y) = \begin{bmatrix} -x + \frac{axy}{(x+y)^2} & -\frac{a(x)^2}{(x+y)^2} \\ \frac{b(y)^2}{(x+y)^2} & -\frac{bxy}{(x+y)^2} \end{bmatrix}. \quad (2.16)$$

The Jacobian matrix evaluated at the boundary equilibrium point  $E_1(1, 0)$  takes the form

$$J_1 = [J(x, y)]_{E_1} = \begin{bmatrix} -1 & -a \\ 0 & b - c \end{bmatrix} \quad (2.17)$$

and therefore, if the positive interior equilibrium point exists (i.e.  $b > c$ ),  $E_1(1, 0)$  is always stable along the  $x$ -direction and unstable along the  $y$ -direction and consequently,  $E_1(1, 0)$  is a saddle point.

At the interior equilibrium point  $E_*(x^*, y^*)$  the Jacobian matrix  $J_*$  is given by

$$J_* = [J(x, y)]_{E_*} = \begin{bmatrix} -x^* + \frac{ax^*y^*}{(x^*+y^*)^2} & -\frac{a(x^*)^2}{(x^*+y^*)^2} \\ \frac{b(y^*)^2}{(x^*+y^*)^2} & -\frac{bx^*y^*}{(x^*+y^*)^2} \end{bmatrix}. \quad (2.18)$$


The characteristic equation for the Jacobian matrix  $J_*$  is given by

$$\lambda^2 + A_1\lambda + A_2 = 0, \quad \text{where } A_1 = -\text{Tr}(J_*) \quad \text{and} \quad A_2 = \det(J_*). \quad (2.19)$$

As  $\det(J_*) = ab(x^*)^2y^*/(x^*+y^*)^2 > 0$  the stability of the equilibrium point  $E_*$  solely depends on the sign of  $\text{Tr}(J_*)$  and hence  $E_*$  is stable if  $\text{Tr}(J_*) < 0$  and unstable if  $\text{Tr}(J_*) > 0$ . It is easy to verify that both the situations are possible for the model system under some appropriate parameter values. In terms of the parameters the stability condition of positive interior equilibrium point  $E_*$  is  $a < b[c + b/(b - c)]/(b + c)$ . We have noted earlier that  $a < b/(b - c)$  and  $b > c$  are the necessary conditions for the positivity of  $E_*$ . Hence the stability of equilibrium point  $E_*$  demands the following condition

$$a < \bar{a} \equiv \min \left[ \frac{b}{b-c}, \frac{b}{b+c} \left( c + \frac{b}{b-c} \right) \right] \quad \text{with } b > c. \quad (2.20)$$





**Figure 1.** The Hopf-bifurcating periodic solution of the model system (2.5) for parametric values  $a = 2.0$ ,  $b = 0.7808$  and  $c = 0.5$ .

If we increase the value of ‘ $a$ ’ further such that  $a > b[c + b/(b - c)]/(b + c)$  with the restriction  $b > c$  then  $E_*$  becomes locally unstable. Hence by the Poincaré criteria [11] there exists at least one limit cycle around  $E_*$  within the positive  $(x, y)$ -plane. We now deduce the condition for the existence of a Hopf-bifurcating small amplitude periodic solution.

**Lemma.** *If  $a = a^* = b[c + b/(b - c)]/(b + c)$  with  $b > c$ , then the system (2.5) exhibits Hopf-bifurcation near  $E_*$ .*

**Proof.** At the parametric value  $a = a^* = b[c + b/(b - c)]/(b + c)$ ,  $\text{Tr}(J_*) = 0$  and  $\det(J_*) > 0$ . When  $a$  takes the value  $a = a^*$ , the roots of the characteristic equation (2.19) are purely imaginary. Also we can verify that the result  $(d/da)[\text{Tr}(J_*)]_{a=a^*} \neq 0$ . Hence both the conditions for Hopf-bifurcation [9, 22, 36, 48] are satisfied (see figure 1).  $\square$

The above result establishes the existence of a small amplitude periodic solution near the interior equilibrium point  $E_*$ . Our next task is to find the condition for stability of the Hopf-bifurcating periodic solution and for this we have to calculate the first Lyapunov coefficient at the critical parametric value  $a = a^*$ . For this purpose we follow the procedure and standard notation as introduced by Kuznetsov [46]. First we translate the origin to the equilibrium point  $E_*$  using the change of variables  $x = x^* + h_1$  and  $y = y^* + h_2$ . Substituting this transformation in (2.5a) and expanding in Taylor series at the critical parametric condition  $a = a^*$

we get

$$\frac{dh_1}{d\tau} = a_{10}h_1 + a_{01}h_2 + a_{20}h_1^2 + a_{11}h_1h_2 + a_{02}h_2^2 + a_{30}h_1^3 + a_{21}h_1^2h_2 + a_{12}h_1h_2^2 + a_{03}h_2^3 + \dots, \quad (2.21a)$$

$$\frac{dh_2}{d\tau} = b_{10}h_1 + b_{01}h_2 + b_{20}h_1^2 + b_{11}h_1h_2 + b_{02}h_2^2 + b_{30}h_1^3 + b_{21}h_1^2h_2 + b_{12}h_1h_2^2 + b_{03}h_2^3 + \dots, \quad (2.21b)$$

where the  $a_{ij}$  and  $b_{ij}$  are given by

$$a_{ij} = \left[ \frac{\partial^{i+j} F_1(x, y)}{\partial x^i \partial y^j} \right]_{a=a^*}, \quad b_{ij} = \left[ \frac{\partial^{i+j} F_2(x, y)}{\partial x^i \partial y^j} \right]_{a=a^*}. \quad (2.22)$$

In the above expansion we are not interested in the coefficients of fourth and higher order terms as they make no contribution to the calculation of the first Lyapunov coefficient. In order to make the coefficients of  $h_1$  in  $dh_1/d\tau$  and of  $h_2$  in  $dh_2/d\tau$  equal to zero we introduce the non-singular transformation of variables as  $h_1 = (a_{10}/b_{10})u_1 + u_2$ ,  $h_2 = u_1$  in the above expansion to get the transformed system as

$$\frac{du_1}{d\tau} = \alpha_{01}u_2 + \alpha_{20}u_1^2 + \alpha_{11}u_1u_2 + \alpha_{02}u_2^2 + \alpha_{30}u_1^3 + \alpha_{21}u_1^2u_2 + \alpha_{12}u_1u_2^2 + \alpha_{03}u_2^3 + \dots, \quad (2.23a)$$

$$\frac{du_2}{d\tau} = \beta_{10}u_1 + \beta_{20}u_1^2 + \beta_{11}u_1u_2 + \beta_{02}u_2^2 + \beta_{30}u_1^3 + \beta_{21}u_1^2u_2 + \beta_{12}u_1u_2^2 + \beta_{03}u_2^3 + \dots. \quad (2.23b)$$

The above system can be written in the following form

$$\frac{d}{d\tau} \mathbf{u} = A\mathbf{u} + \frac{1}{2}B(\mathbf{u}, \mathbf{u}) + \frac{1}{6}C(\mathbf{u}, \mathbf{u}, \mathbf{u}) + \dots, \quad (2.24)$$

where  $A$  denotes the coefficient matrix of linear part with zero as the main diagonal elements;  $B(\mathbf{u}, \mathbf{v})$ ,  $C(\mathbf{u}, \mathbf{v}, \mathbf{w})$  are symmetric multi-linear vector functions of  $\mathbf{u} = (u_1, u_2)$ ,  $\mathbf{v} = (v_1, v_2)$ ,  $\mathbf{w} = (w_1, w_2) \in \mathbf{R}^2$  and take the following forms:

$$B(\mathbf{u}, \mathbf{v}) = \begin{bmatrix} 2\alpha_{20}u_1v_1 + \alpha_{11}(u_1v_2 + u_2v_1) + 2\alpha_{02}u_2v_2 \\ 2\beta_{20}u_1v_1 + \beta_{11}(u_1v_2 + u_2v_1) + 2\beta_{02}u_2v_2 \end{bmatrix}, \quad (2.25)$$

$$C(\mathbf{u}, \mathbf{v}, \mathbf{w}) = \begin{bmatrix} c_{11} + c_{12} + c_{13} + c_{14} \\ c_{21} + c_{22} + c_{23} + c_{24} \end{bmatrix}, \quad (2.26)$$

where the  $c_{ij}$  ( $i = 1, 2$ ;  $j = 1, 2, 3, 4$ ) are given by

$$c_{11} = 6\alpha_{30}u_1v_1w_1, \quad c_{12} = 2\alpha_{21}(u_1v_1w_2 + u_1v_2w_1 + u_2v_1w_1),$$

$$c_{13} = 2\alpha_{12}(u_1v_2w_2 + u_2v_1w_2 + u_2v_2w_1), \quad c_{14} = 6\alpha_{03}u_2v_2w_2,$$

$$c_{21} = 6\beta_{30}u_1v_1w_1, \quad c_{22} = 2\beta_{21}(u_1v_1w_2 + u_1v_2w_1 + u_2v_1w_1),$$

$$c_{23} = 2\beta_{12}(u_1v_2w_2 + u_2v_1w_2 + u_2v_2w_1), \quad c_{24} = 6\beta_{03}u_2v_2w_2.$$

Let  $\lambda_{1,2} = \pm i\omega$  be the eigenvalues of the matrix  $A$  and  $\mathbf{p}, \mathbf{q}$  proper eigenvectors satisfying the relations

$$A\mathbf{q} = i\omega\mathbf{q}, \quad A^T\mathbf{p} = -i\omega\mathbf{p} \quad \text{and} \quad \langle \mathbf{p}, \mathbf{q} \rangle = 1, \quad (2.27)$$

where  $\langle \cdot, \cdot \rangle$  means the standard scalar product in  $\mathbb{C}^2$ :  $\langle \mathbf{p}, \mathbf{q} \rangle = \bar{p}_1 q_1 + \bar{p}_2 q_2$ . The first Lyapunov coefficient determining the stability of the Hopf-bifurcating periodic solution is given by [46]

$$l_1 = \frac{1}{2\omega^2} \operatorname{Re}(i g_{20} g_{11} + \omega g_{21}). \tag{2.28}$$

The quantities  $g_{20}$ ,  $g_{11}$  and  $g_{21}$  are given by

$$g_{20} = \langle \mathbf{p}, B(\mathbf{q}, \mathbf{q}) \rangle, \quad g_{11} = \langle \mathbf{p}, B(\mathbf{q}, \bar{\mathbf{q}}) \rangle, \quad g_{21} = \langle \mathbf{p}, C(\mathbf{q}, \mathbf{q}, \bar{\mathbf{q}}) \rangle. \tag{2.29}$$

The stability of the Hopf-bifurcating periodic solution depends on the sign of first Lyapunov coefficient. The limit cycle is called stable if  $l_1 < 0$  and is unstable for  $l_1 > 0$ . The explicit expression for  $g_{20}$ ,  $g_{11}$ ,  $g_{21}$  and hence that of the Lyapunov coefficient  $l_1$  in terms of the system parameters is too lengthy and would take at least two printed pages. Furthermore, it is quite difficult to conclude about the sign of  $l_1$  due to its complicated algebraic expression. For this reason we present here a numerical example to illustrate the method of calculation for  $l_1$ . For numerical calculation of the first Lyapunov coefficient  $l_1$ , we consider the following model

$$\frac{dx}{d\tau} = x(1-x) - \frac{2xy}{x+y} \equiv F_1(x, y), \quad \frac{dy}{d\tau} = \frac{0.7807764064xy}{x+y} - 0.5y \equiv F_2(x, y). \tag{2.30}$$

From the previous analysis, one can easily verify that the positive equilibrium point is  $E_* = (0.2807764064, 0.1576707808)$  and the system exhibits limit cycle oscillation at the critical parametric value  $a^* = 2$ . Let  $x = 0.2807764064 + h$ ,  $y = 0.1576707808 + k$ , then system (2.30) becomes

$$\begin{aligned} \frac{dh}{d\tau} = & 0.1798058984h - 0.8201941014k - 0.4100970509h^2 - 2.100970508hk \\ & + 1.870679355k^2 - 1.345436728h^3 + 3.446407236kh^2 \\ & + 0.5252426287k^2h - 4.266601336k^3, \end{aligned} \tag{2.31a}$$

$$\begin{aligned} \frac{dk}{d\tau} = & 0.1009705080h - 0.1798058985k - 0.2302911524h^2 + 0.8201941016hk \\ & - 0.7302911521k^2 + 0.5252426269h^3 - 1.345436728kh^2 \\ & - 0.2050485261k^2h + 1.665630829k^3. \end{aligned} \tag{2.31b}$$

Making the non-singular transformation  $h = (0.1798058983/0.1009705080)u + v$ ,  $k = u$ , we see that system (2.31) becomes


$$\begin{aligned} \frac{du}{d\tau} = & 0.1009705080v - 0.00000001u^2 + 0.000000002vu - 0.2302911524v^2 \\ & - 0.0000001u^3 - 0.0000011vu^2 + 1.460582304v^2u + 0.5252426269v^3, \end{aligned} \tag{2.32a}$$

$$\begin{aligned} \frac{dv}{d\tau} = & -0.499999995u - 3.171164609u^2 - 3.561552813vu - 0.000000002v^2 \\ & + 0.0000013u^3 + 0.0000161vu^2 - 6.342329213v^2u - 2.280776405v^3. \end{aligned} \tag{2.32b}$$

$\lambda_{1,2} = \pm i\omega = \pm 0.2246892386i$  are the eigenvalues of the coefficient matrix corresponding to the linear part of (2.32) where the matrix  $A$  is given by,

$$A = \begin{bmatrix} 0 & 0.1009705080 \\ -0.499999995 & 0 \end{bmatrix}.$$

The eigenvectors as defined in (2.27) are given by  $\mathbf{p} = (0.1754320563, 0.07883539039i)$  and  $\mathbf{q} = (0.07883539039, 0.1754320563i)$ . Now one can calculate the quantities  $g_{20}$ ,  $g_{11}$ ,  $g_{21}$



**Figure 2.** Stable limit cycle around the positive interior equilibrium point for the model system (2.30).

easily by using any mathematical software (e.g. MAPLE) as

$$g_{20} = -0.005\,279\,657\,042 + 0.003\,107\,509\,389i,$$

$$g_{11} = 0.005\,279\,657\,042 + 0.003\,107\,509\,388i,$$

$$g_{21} = -0.004\,581\,433\,107 + 0.005\,411\,288\,533i.$$

Hence the first Lyapunov coefficient  $l_1$  is given by

$$l_1 = -0.201\,941\,0159.$$


Thus the Hopf-bifurcating periodic solution is stable and all other trajectories around the limit cycle ultimately approach it (see figure 2).

### 2.5. Global stability

In the previous section we have obtained the conditions for the existence of positive equilibrium and its local asymptotic stability conditions. We have observed that instability of the boundary equilibrium  $E_1$  gives support for the existence of positive interior equilibrium point  $E_*$ . The parametric conditions for local asymptotic stability of  $E_*$  are  $a < \bar{a}$  and  $b > c$ . Now we try to find the condition under which system (2.5) will have no non-trivial periodic solutions around  $E_*$ . For this purpose we recall the divergence criterion for the stability of a periodic solution for planar systems [35, 37].

Let us construct the function  $h(x, y) = 1/(xy)$  such that  $h(x, y) > 0$  for all  $x > 0, y > 0$ . Using the definition of  $F_1(x, y)$  and  $F_2(x, y)$  from equation (2.2) we get

$$\frac{\partial}{\partial x}(F_1 h) + \frac{\partial}{\partial y}(F_2 h) = -\frac{1}{y} - \frac{(b-a)}{(x+y)^2}.$$



**Figure 3.** The phase portrait of the system (2.5) created by MATLAB.  $E_*$  is a global attractor, where  $a = 0.7$ ,  $b = 0.9$  and  $c = 0.8$ .

Clearly

$$\Delta(x, y) \equiv \frac{\partial}{\partial x}(F_1h) + \frac{\partial}{\partial y}(F_2h) < 0 \quad \text{for } a < b.$$

According to the Bendixon–Dulac criterion, there will be no limit cycle in the positive quadrant of the  $xy$ -plane. Now we can state the following lemma.

**Lemma.** *The existence of interior equilibrium point  $E_*$  along with its local stability and the restriction  $a < b$  eliminates the chance of existence of a non-trivial periodic solution around  $E_*$ .*

Now we are in a position to prove the global stability of the model system (2.5).

**Lemma.** *If  $a < a_*$  [ $a_* = \min(b, \bar{a})$ ] and  $b > c$  then  $E_*$  is globally asymptotically stable.*

**Proof.** The system (2.5) has no positive periodic solution around  $E_*$  for  $a < a_*$  and  $b > c$ . The boundedness of the solution together with the saddle nature of boundary equilibrium point  $E_1$  and local asymptoticity of  $E_*$  leads to the conclusion that all the trajectories will approach  $E_*$  with increasing time ‘ $t$ ’. Hence  $E_*$  is a global attractor and the system is globally asymptotically stable (see figure 3).

### 3. The stochastic model

The above discussion rests on the assumption that the environmental parameters involved with the model system are all constants irrespective of time and environmental fluctuations. In reality all such parameters exhibit random variations to a greater or lesser extent. In previous discussions we have dealt with the equilibrium populations and their stability with regard to the imposition of small disturbances. In this section we consider the effect of environmental

fluctuations on the model system and the stochastic stability of the coexisting equilibrium point associated with the model system.

Environmental fluctuations are important components in an ecosystem. Most natural phenomena do not follow strictly deterministic laws but rather oscillate randomly about some average value so that the deterministic equilibrium is no longer an absolutely fixed state [5, 9]. May [49] pointed out the fact that due to environmental fluctuations, the birth rates, carrying capacity, competition coefficients and other parameters involved with the model system exhibit random fluctuations to a greater or lesser extent. Consequently the equilibrium population distribution fluctuates randomly around some average value. Elton [25] observed that 'the chief cause of fluctuations in animal numbers is the instability of the environment'. Within a deterministic environment we seek the constant equilibrium population and then investigate its stability which follows from the dynamics of the interactions between and within the species. For systems which are driven by environmental stochasticity, it is impossible to find a time-independent equilibrium point as a solution of the governing SDEs. In this situation it is reasonable to find a probabilistic 'smoke cloud', described by the equilibrium probability distribution. For the model systems described by the system of SDEs, there is a continuous spectrum of disturbances generated by the environmental stochasticity, and the system is in tension between two countervailing tendencies. On the one hand, random environmental fluctuations are responsible for spreading the cloud and making the probability distribution move diffusively, while on the other hand, the dynamics of stabilizing population interactions tend to restore the populations to their mean value in order to compact the cloud [49]. Model systems with this type of compact cloud of population distribution are called stochastically stable systems. To study the effect of random environmental fluctuations we have to construct the stochastic counterpart of the deterministic model system by incorporating environmental fluctuations.

There are two ways of developing the stochastic model corresponding to an existing deterministic one to study the effect of fluctuating environment. First, one can replace the environmental parameters involved with the deterministic model system by some random parameters (e.g. the growth rate parameter ' $r$ ' can be replaced by  $r_0 + \epsilon \gamma(t)$ , where  $r_0$  is the average growth rate,  $\gamma(t)$  is the noise function and  $\epsilon$  is the intensity of fluctuation). Second, one can add a randomly fluctuating driving force directly to the deterministic growth equations of prey and predator populations without altering any particular parameter [8, 11, 59].

Model (2.2) was just a first attempt towards the modelling of predator-prey interaction with ratio-dependent functional response. In this study we introduce stochastic perturbation terms into the growth equations of both prey and predator populations to incorporate the effect of randomly fluctuating environment. We assume that stochastic perturbations of the state variables around their steady-state values  $E_*$  are of Gaussian white noise type which are proportional to the distances of  $x, y$  from their steady-state values  $x^*, y^*$ , respectively [15]. Gaussian white noise is extremely useful to model rapidly fluctuating phenomena [11, 59]. So the deterministic model system (2.2) results in the following stochastic model system:

$$dx = F_1(x, y)dt + \sigma_1(x - x^*)d\xi_1^1, \quad dy = F_2(x, y)dt + \sigma_2(y - y^*)d\xi_1^2, \quad (3.1)$$

where  $\sigma_1, \sigma_2$  are real constants and known as the intensity of environmental fluctuations,  $\xi_i^j = \xi_i^j(t), i = 1, 2$  are standard Wiener processes independent of each other [30]. In the rest of this work we consider (3.1) as an Ito stochastic differential system of the type

$$dX_t = f(t, X_t)dt + g(t, X_t)d\xi_t, \quad X_{t_0} = X_0, \quad (3.2)$$

where the solution  $(X_t, t > 0)$  is an Ito process, ' $f$ ' is a slowly varying continuous component or *drift coefficient*, ' $g$ ' is the rapidly varying continuous random component or *diffusion coefficient* and  $\xi_t$  is a two-dimensional stochastic process having scalar Wiener process

components with increments  $\Delta \xi_j^j = \xi_j(t + \Delta t) - \xi_j(t)$  that are independent Gaussian random variables  $N(0, \Delta t)$ . In the case of system (3.1),

$$\begin{aligned} X_t &= (x, y)^T, & \xi_t &= (\xi_t^1, \xi_t^2)^T, & f &= \begin{bmatrix} F_1(x, y) \\ F_2(x, y) \end{bmatrix}, \\ g &= \begin{bmatrix} \sigma_1(x - x^*) & 0 \\ 0 & \sigma_2(y - y^*) \end{bmatrix}. \end{aligned} \tag{3.3}$$

Since the diffusion matrix ‘g’ depends on the solution  $X_t$ , system (3.1) is said to have multiplicative noise.

### 3.1. Stochastic stability of interior equilibrium

The stochastic differential system (3.1) can be centred at its positive equilibrium point  $E_*(x^*, y^*)$  by introducing the variables  $u_1 = x - x^*$  and  $u_2 = y - y^*$ . It looks a very hard problem to derive asymptotic stability in the mean square sense by the Lyapunov function method working on the complete nonlinear equations (3.1). For simplicity of mathematical calculations we deal with the SDEs obtained by linearizing the vector function ‘f’ in (3.3) about the positive equilibrium point  $E_*$ . The linearized version of (3.2) around  $E_*$  is given by

$$dU(t) = F(U(t))dt + g(U(t))d\xi(t), \tag{3.4}$$

where  $U(t) = \text{col}(u_1(t), u_2(t))$  and

$$F(U(t)) = \begin{bmatrix} -a_{11}u_1 - a_{12}u_2 \\ a_{21}u_1 - a_{22}u_2 \end{bmatrix}, \quad g(U(t)) = \begin{bmatrix} \sigma_1 u_1 & 0 \\ 0 & \sigma_2 u_2 \end{bmatrix} \tag{3.5}$$

with

$$\begin{aligned} a_{11} &= x^* - \frac{ax^*y^*}{(x^* + y^*)^2}, & a_{12} &= \frac{a(x^*)^2}{(x^* + y^*)^2}, \\ a_{21} &= \frac{b(y^*)^2}{(x^* + y^*)^2}, & a_{22} &= \frac{bx^*y^*}{(x^* + y^*)^2}. \end{aligned} \tag{3.6}$$

Note that, in (3.4) the positive equilibrium  $E_*$  corresponds to the trivial solution  $(u_1, u_2) = (0, 0)$ . Let  $\Omega$  be the set defined by  $\Omega = \{t \geq t_0\} \times \mathbb{R}^2, t_0 \in \mathbb{R}^+$ . Let  $V \in C_2(\Omega)$  be a twice differentiable function of time  $t$ . We define the following theorem due to Afanas’ev *et al* [2].

**Theorem.** *Suppose there exists a function  $V(U, t) \in C_2(\Omega)$  satisfying the inequalities*

$$K_1|U|^\alpha \leq V(U, t) \leq K_2|U|^\alpha, \tag{3.7}$$

$$LV(U, t) \leq -K_3|U|^\alpha, \quad K_i > 0, \quad i = 1, 2, 3, \quad \alpha > 0. \tag{3.8}$$

*Then the trivial solution of (3.4) is exponentially  $\alpha$ -stable for all time  $t \geq 0$ .*

With reference to (3.8) the expression for  $LV(U, t)$  is defined by

$$LV(U, t) = \frac{\partial V(U, t)}{\partial t} + F^T(U) \frac{\partial V(U, t)}{\partial U} + \frac{1}{2} \text{Tr} \left[ g^T(U) \frac{\partial^2 V(U, t)}{\partial U^2} g(U) \right], \tag{3.9}$$

where

$$\frac{\partial V(U, t)}{\partial U} = \text{col} \left( \frac{\partial V}{\partial u_1}, \frac{\partial V}{\partial u_2} \right), \quad \frac{\partial^2 V(U, t)}{\partial U^2} = \left[ \left( \frac{\partial^2 V}{\partial u_i \partial u_j} \right)_{i,j=1,2} \right]. \tag{3.10}$$

Let us consider the Lyapunov function

$$V(U(t), t) = \frac{1}{2} [u_1^2 + \omega_1 u_2^2], \tag{3.11}$$

where  $\omega_1$  is a positive real constant to be chosen later. It can be easily checked that (3.7) holds for the Lyapunov function defined in (3.11) with  $\alpha = 2$ . Now,

$$LV(U, t) = (-a_{11}u_1 - a_{12}u_2)u_1 + (a_{21}u_1 - a_{22}u_2)\omega_1 u_2 + \frac{1}{2} \text{Tr} \left[ g^T(U) \frac{\partial^2 V(U, t)}{\partial U^2} g(U) \right]. \tag{3.12}$$

From (3.5), (3.10) and (3.11) we get,

$$\frac{\partial^2 V}{\partial u^2} = \begin{bmatrix} 1 & 0 \\ 0 & \omega_1 \end{bmatrix}, \quad g^T(U) \frac{\partial^2 V(U, t)}{\partial U^2} g(U) = \begin{bmatrix} \sigma_1^2 u_1^2 & 0 \\ 0 & \omega_1 \sigma_2^2 u_2^2 \end{bmatrix}. \tag{3.13}$$

Hence from (3.12) we get,

$$LV(U, t) = - \left( 2a_{11} - \frac{\sigma_1^2}{2} \right) u_1^2 + 2(a_{21}\omega_1 - a_{12})u_1 u_2 - \left( 2a_{22} - \frac{\omega_1 \sigma_2^2}{2} \right) u_2^2.$$

If we choose  $\omega_1 = (a_{12}/a_{21}) > 0$ , then from the above result we get,

$$LV(U, t) = - \left( 2a_{11} - \frac{\sigma_1^2}{2} \right) u_1^2 - \left( 2a_{22} - \frac{a_{12}\sigma_2^2}{2a_{21}} \right) u_2^2 = -U^T Q U, \tag{3.14}$$

where  $Q = \text{diag}[(2a_{11} - \sigma_1^2/2), (2a_{22} - a_{12}\sigma_2^2/2a_{21})]$  and the diagonal matrix  $Q$  will be a real symmetric positive definite matrix and hence its eigenvalues  $\lambda_1$  and  $\lambda_2$  will be positive real quantities if and only if the following conditions hold:

$$\sigma_1^2 < 4a_{11} \quad \text{with } a_{11} > 0 \quad \text{and} \quad \sigma_2^2 < \frac{4a_{22}a_{21}}{a_{12}}. \tag{3.15}$$

If  $\lambda_m$  stands for the minimum of two positive eigenvalues  $\lambda_1$  and  $\lambda_2$  for the diagonal matrix  $Q$  then from (3.14) we get the following result:

$$LV(U, t) \leq -\lambda_m |U|^2. \tag{3.16}$$

This leads us to the following theorem.

**Theorem.** Assume that for some positive real value of  $\omega_1 = a_{12}/a_{21}$  and the inequalities in (3.15) hold then the zero solution of system (3.4) is asymptotically mean square stable.

Recall that  $a < \bar{a}$  and  $b > c$  are the conditions for deterministic stability of the interior equilibrium point  $E_*$ . Conditions for deterministic stability of interior equilibrium point along with the inequalities (3.15) are the necessary conditions for stochastic stability of the model system under environmental fluctuation. Inequalities (3.15) defines the upper threshold values for the intensities of the environmental fluctuations ‘ $\sigma_1$ ’ and ‘ $\sigma_2$ ’ determined by the system parameters (i.e.  $a, b$  and  $c$ ) as

$$\sigma_1^2 < \bar{\sigma}_1^2 = 4 \left[ \frac{b^2 - a(b-c)^2}{b^2} \right] \quad \text{and} \quad \sigma_2^2 < \bar{\sigma}_2^2 = \frac{4(b-c)^3}{ac}. \tag{3.17}$$

Thus the internal parameters of the model system and the intensities of environmental fluctuation have the ability to maintain the stability of the stochastic model system and exhibit a balanced dynamics at any future time within a bounded domain of  $(a, b, c, \sigma_1, \sigma_2)$ -parametric space. The boundaries of the bounded set in  $(a, b, c, \sigma_1, \sigma_2)$ -parametric space are defined by the following inequalities (which are some implicit functional relations):

$$a < \bar{a}, \quad b > c, \quad \sigma_1^2 < \bar{\sigma}_1^2, \quad \sigma_2^2 < \bar{\sigma}_2^2, \tag{3.18}$$

where the expressions for  $\bar{a}$ ,  $\bar{\sigma}_1^2$  and  $\bar{\sigma}_2^2$  are given in (2.20) and (3.17), respectively. The inequalities in (3.17) can be put into an alternative form as

$$a < \left[ \frac{b}{b-c} \right]^2 \left[ 1 - \frac{\sigma_1^2}{4} \right] \quad \text{and} \quad a < \frac{4(b-c)^3}{c\sigma_2^2}. \tag{3.19}$$



For a given set of values for  $b, c, \sigma_1$  and  $\sigma_2$  with  $b > c$  we can find an estimate for the parameter ‘ $a$ ’ which will ensure the deterministic stability as well as stochastic stability of interior equilibrium point  $E_*$  for the model system (2.5). Defining the upper threshold limit ‘ $a$ ’ for ‘ $a$ ’ as

$$A = \min \left[ \frac{b}{b-c}, \frac{b}{b+c} \left( c + \frac{b}{b-c} \right), \left[ \frac{b}{b-c} \right]^2 \left[ 1 - \frac{\sigma_1^2}{4} \right], \frac{4(b-c)^3}{c\sigma_2^2} \right] \tag{3.20}$$

we can conclude that  $a < A$  and  $b > c$  are the necessary and sufficient conditions for the stochastic stability of interior equilibrium point  $E_*$  for the model system under consideration.

### 3.2. Numerical simulation

In order to give some support to the stability results of the stochastic model system obtained in the previous section, we numerically simulate the solution of the SDE (3.1). For this purpose we have to keep in mind that approximated sample paths or trajectories of Ito processes obtained from direct simulation must be close to those of the original Ito process and these will lead us to the concept of a strong solution for a system of SDE [21]. To find the approximate strong solution of the Ito system of SDEs (3.1) with given initial condition we use the Euler–Maruyama (EM) and Milstein method.

Consider the discretization of the time interval  $[t_0, t_f]$  with

$$t_0 = 0 < t_1 < t_2 < \dots < t_n < \dots < t_N < t_{N+1} = t_f$$

and the simplest stochastic numerical scheme for the system under consideration is the EM method

$$u_{k,n+1} = u_{k,n} + f(t_n, u_{k,n})\Delta t_n + g(t_n, u_{k,n})\Delta \xi_n^k$$

with  $u_{k,0} = u_{k0}, k = 1, 2$  and  $\bar{u}_n = [u_{1,n}, u_{2,n}]$  being the numerical solution at time ‘ $t_n$ ’. In the above numerical scheme, the increments are given by

$$\begin{aligned} \Delta t_n &= t_{n+1} - t_n, \\ \Delta \xi_n^k &= \xi_{n+1}^k - \xi_n^k = \xi^k(t_{n+1}) - \xi^k(t_n), \end{aligned}$$

where  $n = 0, 1, 2, \dots, N$ .

The noise increments  $\Delta \xi_n^k$  are  $N(0, \Delta t_n)$ -distributed independent random variables which can be generated numerically by pseudo-random number generators.

An efficient way to evaluate the increments of the Wiener process  $\Delta \xi_n^k$  is to consider


$$\Delta \xi_n^k = \sqrt{I_{nk}} \Delta t_n,$$

where  $I_{nk}$  is the  $n$ th realization of  $I_k$  and  $I_k$  is the Gaussian random variable  $N(0, 1)$ .


Figures 4 and 5 are generated by using the numerical simulation method discussed above with the help of MATLAB software.

### 4. Conclusion

In classical predator–prey model systems, the consumption rate of a single predator (so-called functional response) is the key component of predation models as it is considered to determine both the prey death rate and the predator rate of increase. In a continuous-time predator–prey model with prey-dependent functional response it is a classical assumption that predators encounter prey at random and that the response function depends solely on prey abundance. This assumption is true for several cases but is not always appropriate [5].



**Figure 4.** Solution of SDE (3.1) with parametric values  $a = 1.9$ ,  $b = 1.5$ ,  $c = 1.2$  and  $\sigma_1 = \sigma_2 = 0.2$ .




**Figure 5.** Corresponding population distribution around the equilibrium point  $E_* = (0.62, 0.156)$ .


In reality, it is reasonable to assume that the response function depends on the ratio of prey to predator abundance. There are several field and laboratory observations in support of ratio-dependent functional response. There is a sharp difference between the two types of response functions with the variations in prey production or abundance. Ratio-dependent models predict proportional increase of both populations while prey-dependent models predict the benefit of predators only from an increase of prey production and abundance.

In this paper we have considered the deterministic analysis of stability for various equilibrium points of a nonlinear predator-prey system with *ratio-dependent* functional response. First we have proved the boundedness of solutions of the model system under consideration. The origin (i.e.  $E_0(0, 0)$ ) and  $E_1(1, 0)$  are two equilibrium points lying on the  $x$ -axis for all permissible parameter values. However the model system cannot be linearized at  $E_0$  and hence its local stability analysis is not possible by traditional analysis with a Jacobian matrix. To overcome this situation we have employed the technique introduced by Arino *et al* [7]. Our analysis shows that the trajectories reach the origin and both the populations become extinct. This may happen in two ways. In one case, absence of prey or predator species is responsible for extinction of predator and prey species, respectively. In the other case, overpredation of prey species irrespective of the growth rate of prey drives the system towards total extinction of both prey and predator species. This result agrees well with the result of Kuang [42]. Next we have considered the local asymptotic stability of interior equilibrium point  $E_*$  and established the existence of a small-amplitude periodic solution arising from Hopf-bifurcation as the parameter ' $a$ ' passes through its critical value ' $a = a^*$ ' from lower to higher. At this position we can conclude that if the consumption ability of the predator is not very high (i.e.  $a < a^*$ ) then both prey and predator populations coexist at their steady-state value  $E_*$  and hence the consumption ability and growing ability generate a balanced dynamics for both the populations. Global stability results are obtained from the condition for non-existence of a trivial periodic solution around  $E_*$  with the parametric restrictions obtained in the last part of section 2.


On the other hand, for the stochastic version of the model system we have obtained the condition for asymptotic stability of equilibrium point  $E_*$  in the mean square sense by using a suitable Lyapunov function (3.11). These conditions depend on  $\sigma_1$ ,  $\sigma_2$  and the parameters involved with the model system. For the deterministic environment, the stability of the equilibrium point demands that all eigenvalues of the Jacobian matrix lie in the left-hand half of the complex plane. For the corresponding model within the stochastic environment, this condition is necessary but insufficient, due to the existence of a relatively compact equilibrium probability cloud for the populations around the deterministic equilibrium point. The stochastic stability requires that the stability provided by the interactions (which is measured by the real parts of eigenvalues of Jacobian matrix) be sufficient to counteract the driving arising from random environmental fluctuations [49]. Regarding stability and instability of the stochastic model system, it intuitively seems appropriate to refer to the systems characterized by large fluctuations in the population numbers as 'unstable' and to those with relatively small fluctuations as 'stable'. For stochastic model system (3.1) asymptotic stability of  $E_*$  in the mean square sense depends on the restriction of (3.15). Recall that the feasible values of the intensities of environmental fluctuations depend on the system parameters, which in turn decrease with the increase of parameter ' $a$ '. For a given set of values of  $a$ ,  $b$  and  $c$  one can easily calculate the upper bounds  $\bar{\sigma}_1^2$  and  $\bar{\sigma}_2^2$  from the relation (3.17). Within the natural environment it is not possible to control the surroundings in such a way that the intensities of environmental fluctuations cannot exceed the upper bounds settled for them by the system parameters. The restrictions (3.15) or equivalently (3.17) are the boundaries determined by the mathematical methods to obtain a stable population distribution around the equilibrium point



**Figure 6.** Solution of SDE (3.1) with parametric values  $a = 1.9$ ,  $b = 1.5$ ,  $c = 1.2$  and  $\sigma_1 = 1.2$  and  $\sigma_2 = 0.8$ .



**Figure 7.** Corresponding population distribution around the equilibrium point  $E_0 = (0.62, 0.156)$ .



**Figure 8.** Solution of SDE (3.1) with parametric values  $a = 1.9$ ,  $b = 1.5$ ,  $c = 1.2$  and  $\sigma_1 = 1.9$  and  $\sigma_2 = 1.8$ .

within a fluctuating environment. Hence we conclude that to preserve the system stochastically stable the above restriction should be maintained.

Now, we are in a position to substantiate the analytical findings for the stochastic model system through numerical experiments. The behaviour of the ecological system depends on the rapidity of environmental fluctuations. This rapidity of fluctuations can be measured through the intensity of fluctuations which act as an environmental driving force. The high amplitude environmental driving force has the ability to change the dynamics of the system from a stable situation to an unstable one. From this viewpoint we have only varied the intensity parameters  $\sigma_1$  and  $\sigma_2$ , keeping all other parameters unaltered. For the numerical analysis, we choose the hypothetical set of parameter values  $a = 1.9$ ,  $b = 1.5$  and  $c = 1.2$  and substituting these values in (2.6) we obtain  $E_* = E_*(0.62, 0.156)$ . Now if we imagine a circular or elliptic neighbourhood around the equilibrium point  $E_*(0.62, 0.156)$ , we find that 90% or more of the population distribution will lie within the said neighbourhood implying the stochastic stability of the model system (see figure 5). If we increase the intensities of environmental fluctuations from  $\sigma_1 = \sigma_2 = 0.2$  to  $\sigma_1 = 1.2$  and  $\sigma_2 = 0.8$  with the same parameter values for  $a$ ,  $b$  and  $c$ , we again find a dense assemblage of probability cloud around  $E_*(0.62, 0.156)$  depicting stochastic stability of the model system (see figures 6 and 7). Now, we would like to see how the behaviour of the system changes for increasing fluctuations. If we further increase  $\sigma_1$  and  $\sigma_2$  (to  $\sigma_1 = 1.9$  and  $\sigma_2 = 1.8$ ) we observe that the amplitude of the fluctuations increases, implying instability of the coexisting equilibrium point within the fluctuating environment (see figure 8). Thus the intensity of the random environmental driving force may be used as

a control parameter. These findings agree well with some earlier works [8, 11, 12, 21, 28, 61]. Numerical simulation establishes the fact that the steady state of the stochastic version is not absolutely fixed but a 'fuzzy' value around which the population fluctuates.

Ecological systems are so complex that environmental fluctuations may not always be controlled. In such a situation, the condition given in (3.20) may provide some feasible way for controlling the species from extinction. It is clear from (3.19) that  $\sigma_1$  never exceeds the value 2.0 as the parameter  $a$  is positive. Hence for a given set of values of  $b$ ,  $c$ ,  $\sigma_1$  and  $\sigma_2$  (with obvious restrictions  $b > c$  and  $0 < \sigma_1 < 2$ ) if we can employ a mechanism to control the predator-prey system such that the parameter value ' $a$ ' lies below the numerical value ' $A$ ' obtained from (3.20) then the ratio-dependent predator-prey model system will exhibit stable fluctuation of population distribution around its interior equilibrium point  $E_*$  embedded within a randomly fluctuating environment.

### Acknowledgments

The authors are grateful to the learned referees for their valuable suggestions which have helped in a better exposition of the paper.

### References

- [1] Abrams P A and Walter C J 1996 Invulnerable prey and the paradox of enrichment *Ecology* **77** 1125–33
- [2] Afanas'ev V N, Kolmanowskii V B and Nosov V R 1996 *Mathematical Theory of Control Systems Design* (Dordrecht: Kluwer)
- [3] Akçakaya H R 1992 Population cycles of mammals: evidence for a ratio-dependent predator-prey hypothesis *Ecol. Monogr.* **62** 119–42
- [4] Arditi R and Berryman A A 1991 The biological control paradox *Trends Ecol. Evol.* **6** 32–43
- [5] Arditi R and Ginzburg L R 1989 Coupling in predator-prey dynamics: ratio-dependence *J. Theor. Biol.* **139** 311–26
- [6] Arditi R, Ginzburg L R and Akçakaya H R 1991 Variation in plankton densities among lakes: a case for ratio-dependent predation models *Am. Nat.* **138** 1287–96
- [7] Arino O, Elabdllaoui A, Mikaram J and Chattopadhyay J 2004 Infection on prey population may act as a biological control in ratio-dependent predator-prey model *Nonlinearity* **17** 1101–16
- [8] Baishya M C and Chakrabarti C G 1987 Non-equilibrium fluctuation in Lotka-Volterra system *Bull. Math. Biol.* **49** 125–31
- [9] Bandyopadhyay M, Bhattacharya R and Chakrabarti C G 2003 A nonlinear two species oscillatory system: bifurcation and stability analysis *Int. J. Math. Math. Sci.* **2003** 1981–91
- [10] Bandyopadhyay M, Bhattacharya R and Mukhopadhyay B 2004 Dynamics of an autotroph-herbivore ecosystem with nutrient recycling *Ecol. Modelling* **146** 201–9
- [11] Bandyopadhyay M and Chakrabarti C G 2003 Deterministic and stochastic analysis of a nonlinear prey-predator system *J. Biol. Syst.* **11** 161–72
- [12] Banerjee S and Chakrabarti C G 1998 Stochastic dynamic modelling of damped Lotka-Volterra system *Syst. Anal. Model. Simul.* **30** 1–10
- [13] Beddington J R, Hassell M P and Lawton J H 1976 The components of arthropod predation—II, the predator rate of increase *J. Anim. Ecol.* **44** 331–40
- [14] Beltrami E and Carroll T O 1995 Modelling the role of viral disease in recurrent phytoplankton blooms *J. Math. Biol.* **32** 857–63
- [15] Beretta E, Kolmanowskii V B and Shaikhet L 1998 Stability of epidemic model with time delays influenced by stochastic perturbations *Math. Comput. Simul.* **45** 269–77
- [16] Beretta E and Kuang Y 1998 Global analysis in some delayed ratio-dependent predator-prey systems *Nonlinear Anal.* **32** 381–408
- [17] Berezovskaya F, Karev G and Arditi R 2001 Parametric analysis of the ratio-dependent predator-prey model *J. Math. Biol.* **43** 221–46
- [18] Berryman A A 1992 The origin and evolution of predator-prey theory *Ecology* **73** 1530–5
- [19] Birkhoff G and Rota G C 1982 *Ordinary Differential Equations* (Boston, MA: Wiley)

- [20] Brezis H 1983 *Analyse Fonctionnelle, Theorie et Applications* (Paris: Masson)
- [21] Carletti M 2002 On the stability properties of a stochastic model for phage–bacteria interaction in open marine environment *Math. Biosci.* **175** 117–31
- [22] Chattopadhyay J and Arino O 1999 A predator–prey model with disease in prey *Nonlinear Anal.* **36** 747–66
- [23] Chattopadhyay J and Bairagi N 2001 Pelican is at risk in Salton sea—an eco-epidemiological model study *Ecol. Modelling* **136** 103–12
- [24] Coe M J, Cumming D H and Phillipson J 1976 Biomass and production of large African herbivores in relation to rainfall and primary production *Oecologia* **22** 341–54
- [25] Elton C S 1966 *The Pattern of Animal Communities* (London: Methuen)
- [26] Freedman H I 1980 *Deterministic Mathematical Models in Population Ecology* (New York: Dekker)
- [27] Freedman H I and Mathsen A M 1993 Persistence in predator–prey systems with ratio-dependent predator influence *Bull. Math. Biol.* **55** 817–27
- [28] Gard T C 1987 *Introduction to Stochastic Differential Equations* (New York: Dekker)
- [29] Gardiner C W 1983 *Handbook of Stochastic Methods* (New York: Springer)
- [30] Gikhman I I and Skorokhod A V 1979 *The Theory of Stochastic Process—I* (Berlin: Springer)
- [31] Gurney W S C and Nisbet R M 1998 *Ecological Dynamics* (USA: Oxford University Press)
- [32] Gutierrez A P 1992 The physiological basis of ratio-dependent predator–prey theory: a metabolic pool model of Nicholson’s blowflies as an example *Ecology* **73** 1552–63
- [33] Haderl K P and Freedman H I 1989 Predator–prey population with parasite infection *J. Math. Biol.* **27** 609–31
- [34] Hairston N G, Smith F E and Slobodkin L B 1960 Community structure, population control and competition *Am. Nat.* **94** 421–7
- [35] Hale J K 1980 *Ordinary Differential Equations* (Malabar, FL: Krieger)
- [36] Hassard B D, Kazarinoff N D and Wan Y H 1981 *Theory and Application of Hopf-Bifurcation* (Cambridge: Cambridge University Press)
- [37] Hsu S B 1978 On the global stability of a predator–prey system *Math. Biosci.* **39** 1–10
- [38] Hsu S B, Hwang T W and Kuang Y 2001 Global analysis of the Michaelis–Menten type ratio-dependent predator–prey system *J. Math. Biol.* **42** 489–506
- [39] Jost C, Arino O and Arditi R 1999 About deterministic extinction in ratio-dependent predator–prey model *Bull. Math. Biol.* **61** 19–32
- [40] Jumarie G 1995 A practical variational approach to stochastic optimal control via state moment equations *J. Frank. Inst. Eng. Appl. Math.* **332** 761–72
- [41] Jumarie G 2003 Stochastics of order in biological systems: applications to population dynamics, thermodynamics, non-equilibrium phase and complexity *J. Biol. Syst.* **11** 113–38
- [42] Kuang Y 1999 Rich dynamics of Gause-type ratio-dependent predator prey system *Fields Inst. Commun.* **21** 325–37
- [43] Kuang Y 2002 Basic properties of mathematical population models *J. Biomath.* **17** 129–42
- [44] Kuang Y and Beretta E 1998 Global qualitative analysis of a ratio-dependent predator–prey system *J. Math. Biol.* **36** 389–406
- [45] Kot M 2001 *Elements of Mathematical Biology* (Cambridge: Cambridge University Press)
- [46] Kuznetsov Y A 1997 *Elements of Applied Bifurcation Theory* (Berlin: Springer)
- [47] Luck R F 1990 Evaluation of natural enemies for biological control: a behavioural approach *Trends. Ecol. Evol.* **5** 196–9
- [48] Marsden J and McCracken M 1976 *The Hopf Bifurcation and its Applications* (New York: Springer)
- [49] May R M 2001 *Stability and Complexity in Model Ecosystems* (Princeton, NJ: Princeton University Press)
- [50] Murray J D 1993 *Mathematical Biology* (New York: Springer)
- [51] Nisbet R M and Gurney W S C 1982 *Modelling Fluctuating Populations* (New York: Wiley Interscience)
- [52] Oksanen T, Oksanen L, Jedrzejewska B, Korpimäki E and Norrdahl K 2000 Predation and the dynamics of the bank vole, *Clethrionomys glareolus* Bank Vole *Biology: Recent Advances in the Population Biology of a Model Species* ed L. Hansson and G Bujalska *Polish J. Ecol.* **48** 197–217
- [53] Oksanen T, Oksanen L, Schneider M and Aunapuu M 2001 Regulation cycles and stability in northern carnivore–herbivore systems: back to first principles *Oikos* **94** 101–17
- [54] Oksanen T, Power M E and Oksanen L 1995 Habitat selection and consumer resources *Am. Nat.* **146** 565–83
- [55] Oksanen T, Schneider M, Rammul Ü, Hambäck P A and Aunapuu M 1999 Population fluctuations of voles in North Fennoscandian tundra: contrasting dynamics in adjacent areas with different habitat composition *Oikos* **86** 463–78
- [56] Rosenzweig M L 1969 Paradox of enrichment: destabilization of exploitation systems in ecological time *Science* **171** 385–7

- [57] Samanta G P 1996 Influence of environmental noises on the Gomati model of interacting species *Ecol. Modelling* **91** 283–91
- [58] Slobodkin L B 1986 The role of minimalism in art and science *Am. Nat.* **127** 252–65
- [59] Tapaswi P K and Mukhopadhyay A 1999 Effects of environmental fluctuation on plankton allelopathy *J. Math. Biol.* **39** 39–58
- [60] Turchin P, Oksanen L, Ekeholm P, Oksanen T and Henttonen H 2000 Are lemmings prey or predators? *Nature* **405** 562–4
- [61] Turelli M 1986 Stochastic community theory: a partially guided tour *Mathematical Ecology* ed T G Hallam and S Levin (Berlin: Springer)
- [62] Venturino E 1995 Epidemics in predator–prey models: disease in the prey *Mathematical Population Dynamics: Analysis of Heterogeneity* vol 1, ed O Arino *et al* (Holland: Wuerz) pp 381–99
- [63] Xiao D and Ruan S 2001 Global dynamics of a ratio-dependent predator–prey system *J. Math. Biol.* **43** 221–90