

# Dynamically recrystallised quartz *c*-axis fabrics in greenschist facies quartzites, Singhbhum shear zone and its footwall, eastern India—influence of high fluid activity

Sojen Joy<sup>1</sup>, Dilip Saha\*

*Geological Studies Unit, Indian Statistical Institute, 203 Barrackpore Trunk Road, Calcutta, 700 035, India*

Received 28 May 1997; accepted 13 January 2000

---

## Abstract

Oriented specimens of quartz tectonites from the Singhbhum shear zone (SSZ), an important tectonic element in the Precambrian eastern Indian shield, and its footwall have been analysed for *c*-axis fabrics of dynamically recrystallised quartz grains. In individual specimens the fabric pattern is either an asymmetric type I cross-girdle or an asymmetric, kinked single-girdle, irrespective of whether the fabric is from the SSZ or its footwall. The asymmetry of the fabric confirms a southward thrust movement of the northern block on a northerly dipping shear zone. The measured fabrics are characterised by a concentration near the Y-axis of the finite strain ellipsoid with an equally or subordinately developed concentration near the Z-axis. Although some earlier works suggest the Y-concentration of the quartz *c*-axis fabric as due to the influence of higher temperature (obtained in amphibolite facies), the occurrence of a similar concentration in the quartz *c*-axis fabric from quartz tectonites of the SSZ and its footwall, interpreted to have formed under greenschist facies condition, is explained by formation of the fabric under high fluid activity during deformation.

---

## 1. Introduction

The Singhbhum shear zone (SSZ), originally named the Copper Belt thrust (CBT) by Dunn and Dey (1942), is an important tectonic element in the Precambrian eastern Indian shield (Fig. 1). Dunn and Dey (1942) considered the overturned southern limb of the Singhbhum anticlinorium to be displaced by the CBT; the rocks of the Chaibasa stage (Chaibasa Formation of Sarkar and Saha, 1977) at the core of the Singhbhum anticlinorium were thrust over the supposedly younger Dhanjori Group lying to the south of and below the SSZ. The Dhanjori Group has been considered as older than the Chaibasa Formation (e.g.

Mukhopadhyay, 1976; Bose and Chakraborty, 1981; Sarkar, 1982b, 1984; Bose, 1994), while others have considered the Dhanjori Group as a molasse sequence younger than the northern metaflysch sequence (Chaibasa Formation of Dunn and Dey, 1942; Gall, 1964; Sarkar and Saha, 1977; Sarkar, 1982a; Saha and Ray, 1984). Gupta et al. (1980) considered the Dhanjori Group and the Chaibasa Formation to be contemporaneous, but representative of different facies (see also Virnave et al., 1994). In the absence of any radiometric data from these two rock groups, and their correct interpretation, the stratigraphic relation (in its current understanding) cannot be utilised for supporting or refuting the interpretation that the SSZ is a major thrust. Some workers have also suggested continuity of structures across the SSZ (Mukhopadhyay, 1976). The metaflysch sequence has a poly-metamorphic history (Bhattacharyya and Sanayal, 1988). Therefore, the apparently higher metamorphic grade north of the SSZ should also be considered in the light

---

\* Corresponding author.

*E-mail addresses:* sojenjoy@del6.vsnl.net.in (S. Joy), dsaha@isica-lac.in (D. Saha).

<sup>1</sup> Now at: BHP Minerals India Pvt. Ltd., Tejkunji, Ambavgarh, Udaipur 313001, India.

of the time relationship between the deformation and metamorphism in the shear zone, the Chaibasa Formation and the Dhanjori Group.

Apart from the obvious copper and uranium mineralisation, the SSZ is marked by the development of a strong foliation and lineation plunging down dip in the foliation plane (cf. *L-S* fabric in mylonites) in rocks varying from pebbly schist (conglomerate), quartzite, quartz–mica schist, to feldspathic chlorite–biotite schist. The strong *L-S* fabric is interpreted to have been induced by strong shearing along the zone (Gall, 1964; Ghosh and Sengupta 1987a,b). For clarity we restrict the use of the term SSZ to the belt of rocks with a strong *L-S* fabric and characterised by copper, uranium, magnetite, apatite and tourmaline mineralisation. The outcrop width of the SSZ passing through the copper and uranium mines of Bhatin, Jaduguda, Rakha Copper Project (Royam), Kendadih, Surda and Musaboni, varies from 500 m to 1 km (Fig. 2).

Development of distinct microstructures and crystallographic preferred orientation (CPO) in minerals including quartz, olivine and calcite have been reported from many large-scale shear zones/thrust

zones (e.g. the Moine thrust zone of Scotland). The microstructures and CPO of minerals in such shear zones have been shown to be useful in understanding the kinematics of deformation, as well as in deducing the ambient pressure–temperature condition (Lister, 1977; Lister and Williams, 1979, 1983; White et al., 1980; Lister and Snoke, 1984; Simpson and Schmid, 1983; Law et al., 1984, 1986; Law, 1987; Jessell, 1988a,b; Wenk et al., 1989; Jessell and Lister, 1990; Gleason et al., 1993). Although the SSZ is known to be a major tectonic element in the Eastern Indian shield, there are few detailed microstructural and fabric studies (Banerji, 1959; Sarkar, 1966, 1984; Sarkar and Bhattacharyya, 1978). The present study documents the SSZ fabrics with a view to comparing them with fabrics from other well-documented crustal-scale shear zones. It also considers the development of deformation-induced fabrics in the Dhanjori quartzites lying below the SSZ and thus considers their evolution in relation to the general deformation in the SSZ. The observed patterns of CPO are interpreted in the light of presently available models of CPO development.




Fig. 1. Geological map of the Singhbhum region (after Dunn and Dey, 1942) with study area outlined. SSZ = Singhbhum shear zone. The Singhbhum shear zone and the Dalma thrust are marked with thick dashed lines.

## 2. Geological background

The study area (Fig. 1) comprises approximately 60 km<sup>2</sup> and covers the central part of the SSZ. The SSZ is a tectonic dislocation zone which separates the Dhanjori Group (footwall) from the Chaibasa Formation (hanging wall) and is marked by intense mylonitisation and formation of a strong *L–S* fabric in the rock units. The mesoscopic structures from the SSZ indicate that the SSZ is a major northerly (or northeasterly) dipping shear zone with a southward thrust movement (Ghosh and Sengupta, 1987a; Joy, 1996). Even within the Dhanjori Group outcrop there is zonal development of strong *L–S* fabric in quartzites, grits and metabasics (a) close to the contact between lower quartzite and metabasic rock unit and (b) in the gritty quartzites and arkoses immediately overlying the Singhbhum Granite (Joy, 1996). These zones are named here as Jublatola shear zone (JSZ) and Rohinbera shear zone (RSZ), respectively (Fig. 2). Singhbhum Granite outcrops along the southern boundary of the study area.

Rocks occurring north of the SSZ within its hanging wall (Fig. 2) contain muscovite–biotite–quartz–almandine as the prevailing metamorphic mineral assemblage (with retrogression of almandine to chlorite). The Dhanjori Group rocks contain chlorite–muscovite–biotite as metamorphic minerals. Rocks within the SSZ usually have the same paragenesis as those in the

Dhanjori Group, except for the local occurrence of kyanite.

## 3. Fabric measurement procedure

Fifty-nine oriented quartz tectonite specimens were collected from the SSZ, the Dhanjori Group south of the SSZ, and from the Chaibasa Formation immediately north of the SSZ in the Singhbhum District, Bihar, India (Fig. 2). The specimens exhibited extensive grain-size reduction by dynamic recrystallisation with the development of a strong crystallographic preferred orientation in the recrystallised quartz (Saha and Joy, 1995; Joy, 1996).

*c*-axis orientations of 200 or more recrystallised quartz grains from each quartzite specimen from the SSZ, and its hanging wall and footwall were measured on sections cut parallel to the mineral elongation lineation and perpendicular to the mesoscopic foliation (XZ section) using a Federov Stage assembly fitted to an optical microscope. A repeat set of measurements was obtained from sections cut perpendicular to both foliation and mineral elongation lineation (YZ section) to check the reproducibility of the fabric (Appendix A). Measurements of the *c*-axis orientations of the relict quartz grains were also carried out where they formed a high proportion of the grains.

The *c*-axis orientations were plotted with respect to




Fig. 2. Location map for specimens collected from the Singhbhum shear zone and the Dhanjori Group.

the mesoscopic foliation–mineral elongation lineation framework on a lower hemisphere equal area projection and contoured using a computer program of Kutty and Joy (1997) to obtain the fabric diagrams presented in this paper. The mineral elongation linea-

tion direction is taken as the principal extension direction (X) of the finite strain ellipsoid. Fabric skeletons were prepared from the contoured fabric diagrams by connecting the crests and ridges (Lister and Williams, 1979; Lister and Hobbs, 1980; Vissers, 1993).




Fig. 3. Micro-structures indicative of high fluid pressure in the Singhbhum shear zone and its footwall. (a) Tuttle lamellae appearing as fine parallel dust inclusion trails in relict and some recrystallised quartz grains, specimen no. rc07, photomicrograph under cross polars. (b) Asymmetric boudin in quartz veins emplaced sub-parallel to mylonitic foliation in the Singhbhum shear zone (white arrows at boudin necks indicate apparent top-to-the-south displacement comparable to that on  $C'$  shears of Passchier and Trouw, 1996). Field photo on a NNE joint surface perpendicular to mylonitic foliation.




Fig. 4. Quartz *c*-axis fabric of the Singhbhum shear zone specimens. Lower hemisphere equal area projections for *c*-axis orientations in dynamically recrystallised grains, *N* is the total number of *c*-axes measured. Fabric patterns are contoured using a computer program of Kuty and Joy (1997) following the Kamb method with  $E = 3\sigma$ . Lowest contour in all fabric diagram corresponds to  $(E-2\sigma)$ . CI is contour interval. The foliation trace is E–W, and the arrow is pointing in the down plunge of mineral elongation lineation direction (taken here as the X direction of the finite strain ellipsoid). Viewing towards east or south-east. Top of foliation is towards north side of the foliation trace. A similar convention is followed for Figs. 5–9. The specimen number is at the top left corner of each fabric diagram. Specimen number rp2 is a quartzite pebble from the Rakha conglomerate and specimen number rc14 is matrix of the Rakha conglomerate.

#### 4. Petrography of the specimens

Quartz tectonites analysed from the study area are composed of quartz as the main constituent with phyllosilicates (white mica and/or chlorite) varying from 2 to 35 modal percent. The quartz tectonites are generally bimodal aggregates with relatively smaller dynamically recrystallised grains (40–120  $\mu\text{m}$ ) and larger relict quartz grains (110–425  $\mu\text{m}$ ). The proportion of the relict grains is typically higher in specimens from the Dhanjori Group (footwall); however, specimens with near to complete recrystallisation are noted from the JSZ also. The relict grains show undulose extinction, deformation bands, subgrain structures, and core-and-mantle structures (White, 1976) and sometimes deformation lamellae (Fairbairn lamellae). The recrystallised quartz grains generally show serrated or sutured grain boundaries, castellate microstructures and, rarely, dragging microstructures (Jessell, 1987). Some of the recrystallised quartz grains also show undulose extinction and incipient subgrain structures (Joy, 1996). A few samples from the Dhanjori Group also show Tuttle lamellae (Tuttle, 1949), which are healed microfractures remaining visible as planes of solid and/or fluid inclusions (Fig. 3a). Quartz veins occurring sub-parallel to mylonitic foliation in the SSZ, but retaining their coarse texture and internal fractures, often show asymmetric boudins with apparent displacement at boudin necks comparable to  $C'$  shears of Passchier and Trouw (1996) (Fig. 3b).

The dynamic recrystallisation is apparently controlled by a combination of both progressive grain-boundary-rotation and grain-boundary-migration recrystallisation. Microstructures indicative of progressive grain boundary rotation—core-and-mantle structure (White, 1976), grain polygonisation (Nicholas and Poirier, 1976); and microstructures indicative of grain-boundary-migration recrystallisation—grain boundary bulging (White, 1976; Bell and Etheridge, 1976), castellate microstructures, dragging microstructures, serrate grain boundaries (Jessell, 1987), and small strain-free grains (White, 1976), are observed in the analysed specimens, either individually or in combination within the same specimen.

#### 5. Description of the quartz $c$ -axis fabric

##### 5.1. Specimens from the Singbhum shear zone

Individual specimens collected from the SSZ are characterised by either of two patterns: (1) an asymmetric, kinked single-girdle pattern oriented at a high angle to the trace of foliation and mineral elongation lineation, with or without the vestige of the trailing edge of a type I cross-girdle, as exemplified by 7m4

and rm15 (Fig. 4) or (2) an asymmetric type I cross-girdle, exemplified by 7m8 (Fig. 4, definitions of fabric types after Lister, 1977). A few specimens display more discontinuous girdles, especially the ones with high mica content, but the connected fabric skeleton of the specimens remains asymmetric (7m1, 11693, Fig. 4).

The fabric point maxima have several locations: close to Y (p593, Fig. 4), close to XZ plane at an angle less than  $45^\circ$  to Z, the foliation normal (e.g. rm11, Fig. 4), or at nearly  $45^\circ$  from Y and Z (e.g. 7m10, Fig. 4). All three types of fabric maxima are equally well developed in some of the analysed specimens (e.g. rm15 and rp2, Fig. 4). Other specimens generally display one or two maxima with or without a submaximum at the other positions described. The central segment of the type I fabric skeleton is oriented at a high angle ( $80$ – $85^\circ$ ) to the foliation trace (e.g. rm15, Fig. 4).

##### 5.2. Specimens from the footwall and hanging wall

Specimens with  $L$ – $S$  fabrics from the footwall display  $c$ -axis fabrics similar to those obtained from the SSZ specimens, except for the details of internal asymmetry and locations of point maxima within the girdle. The specimens from the upper quartzite (Fig. 2) are characterised by either an asymmetric, kinked single-girdle (rm100 and rm99a, Fig. 5), or an asymmetric type I cross-girdle (21093 and rc11, Fig. 5). The point maximum within the girdle is oriented either near Y (rm100, Fig. 5), or close to XZ plane at low angle ( $< 45^\circ$ ) to Z (rc09, Fig. 5), or occasionally at an intermediate orientation (rc13, Fig. 5). Specimens from the JSZ display more strongly developed single-girdle fabrics (9293, rc24, rc03, and 33293, Fig. 6), which are morphologically close to those from the SSZ. The main fabric type in specimens away from the lower quartzite–metabasic rock contact, i.e. the JSZ (Fig. 2), is an asymmetric type I cross-girdle (rc07, Fig. 7). Occasionally fabrics in the lower quartzite away from the JSZ are nearly symmetric type I cross-girdle fabrics (rc37, Fig. 7). A few specimens from the lower quartzite display a more diffused fabric (e.g. rc06 and rc36, Fig. 7), although strongly developed single girdle fabric is also recorded from the lower quartzite (e.g. 4393 and 4993 of Fig. 7).

Only two specimens were analysed for  $c$ -axis fabric from the hanging wall of the SSZ, both specimens containing a very high proportion of mica. One of them (3m3, Fig. 8) displays a poorly developed (type I?) girdle, and the other (4m2, Fig. 8), displays a girdle fabric with a slightly asymmetric central segment, but with point maxima both at Y and at low angle to Z.

## 6. *c*-Axis fabric in the relict grains

With progressive plastic deformation and dynamic recrystallisation, quartzite evolves into a rock with bimodal grain-size distribution consisting of recrystallised quartz and relict porphyroclasts (Marjoribanks, 1976; White, 1976). Krischner and Teysier (1991) have suggested that the CPO in the relict grains would be influenced by (and thus indicative of) the kinematic framework present during most of the crystal-plastic deformation history while the fabric of the recrystallised aggregates reflects the last stages of plastic deformation.

To gain information on the variations, if any, between the *c*-axis fabric of the recrystallised and relict quartz grains, a few specimens from SSZ with rela-

tively high proportions of relict quartz grains were selected for measurement of *c*-axes in relict quartz grains. Where the relict porphyroclasts display undulose extinction, the vector mean of measurements from different parts of the same grain was taken as the average orientation of the relict porphyroclast. The measured fabrics do not indicate any noticeable variation from the fabric of the recrystallised quartz grains (Fig. 9).

## 7. Discussion

The *c*-axis fabric from Singhbhum with the central segment of the fabric skeleton oriented at a high angle (80–85°) to the mesoscopic foliation may be inter-




Fig. 5. Quartz *c*-axis fabric diagrams for the specimens of 'upper quartzite' of the Dhanjori Group (in the eastern part of the study area the Singhbhum shear zone affects the 'upper quartzite' and thus some specimens like p593 may be considered as belonging to the Singhbhum shear zone and the corresponding fabric is included in Fig. 4).

preted in terms of a progressive simple-shear deformation with high shear strain (Lister and Williams, 1979; Bouchez et al., 1983; Lister and Snoke, 1984; Davis et al., 1987). However, it should be noted that *c*-axis fabrics often record only the last episodes of strain (Lister and Price, 1978; Lister and Williams, 1979; Lister and Hobbs, 1980).

### 7.1. Shear sense from quartz *c*-axis fabric

The external asymmetry (as defined by Law, 1987) in the studied quartz *c*-axis fabrics from the SSZ indicate a north-side-up displacement (Fig. 10) consistent with southward thrust movement along a northerly dipping SSZ. The specimens from the Jublatola shear




Fig. 6. Quartz *c*-axis fabric diagrams of the quartz tectonite specimens from the Jublatola shear zone (Dhanjori Group).



zone, except, 23093, also have external asymmetry which can be interpreted in terms of top-to-south displacement (Fig. 6). The opposite sense of asymmetry displayed by 23093 might be due to local aberration in the flow pattern (Passchier, 1983). Some workers anticipate such heterogeneity of flow in 'real' rocks (Jiang and White, 1995).

### 7.2. CPO, slip systems and ambient deformation temperature

Crystallographic preferred orientation in quartz aggregates is an indicator of deformation temperature since the fabric pattern is controlled by the relative importance of different slip systems, which in turn are




Fig. 7. Quartz *c*-axis fabric diagrams of the 'lower quartzite' specimens of the Dhanjori Group.

temperature sensitive. Slip in quartz occurs primarily in the  $\langle a \rangle$  crystallographic direction, predominantly on the basal and prism planes (Schmid and Casey, 1986; Ralser et al., 1991). Basal  $\langle a \rangle$  slip is dominant at lower temperatures and at faster strain rates, causing a  $c$ -axis fabric maximum near the Z-axis of the finite strain ellipsoid. With increasing temperature, rhombohedral  $\langle a \rangle$  slip system becomes activated, causing a fabric maximum at an intermediate orientation between the Y- and Z-axes. Ultimately, at still higher temperature and slower strain rates, the prism  $\langle a \rangle$  slip system operates, causing a fabric maximum near the Y-axis of the finite strain ellipsoid (Tullis et al., 1973; Nicholas and Poirier, 1976; Bouchez, 1977; Tullis, 1977; Lister and Dornsiepen, 1982; Hobbs, 1985; Mainprice et al., 1986; Schmid and Casey, 1986; Culshaw, 1987; Wenk et al., 1989; Jessell and Lister, 1990). Experimental data indicate that the transition temperature from basal  $\langle a \rangle$  to prism  $\langle a \rangle$  slip occurs at about 700–800°C, under a strain rate of  $10^{-5} \text{ s}^{-1}$  (Blacic, 1975). Prism  $\langle c \rangle$  slip system may be activated at very high temperatures and causes a  $c$ -axis fabric concentration near the X direction (Blacic, 1975; Garbutt and Teyssier, 1991). High temperature (700–800°C) and a high partial pressure of water is invoked to explain the activation of prism  $\langle c \rangle$  slip (Blumenfeld et al., 1986; Duebendorfer and Houston, 1987). Quartz  $c$ -axis fabrics with girdles through Y and containing a Y-maximum have been reported from naturally deformed quartzites under amphibolite facies condition (Bunge and Wenk, 1977; Schmid and Casey, 1986), and from granites deformed under subsolidus condition at a temperature of 650–700°C (Blumenfeld et al., 1986). However, Lister and Dornsiepen (1982) recorded notable  $c$ -axis maxima at the Y-axis in quartzites deformed in greenschist to mid-amphibolite facies condition from Saxony granulite terrain.

Specimens from the SSZ and in its footwall are characterised by either asymmetric, kinked single-girdle

or asymmetric type I cross-girdle fabrics passing through Y, at a high angle to the foliation. Often the fabric girdles contain a strong Y-maximum together with equally well developed maxima or submaxima at low angles to Z or at an intermediate orientation in the girdle (Fig. 11).

The metamorphic mineral paragenesis in the Ghatsila–Galudih region indicates that the garnet zone rocks of Ghatsila are replaced southward (still in the hanging wall of the SSZ) by staurolite–kyanite zone rocks (Naha, 1965). According to Naha (1965) these staurolite–kyanite zone rocks (supposedly of amphibolite facies) have suffered retrogression along the SSZ due to dislocation metamorphism. However, Ghosh and Sengupta (1987b) interpreted the development of kyanite as broadly synchronous with progressive deformation in the SSZ. One could explain the Y-maximum as due to supposedly amphibolite facies condition during deformation in the SSZ. However, the Y-maximum fabrics are also present in the JSZ and the lower quartzite in the footwall (Figs. 5–7), where the mineral paragenesis of quartz–chlorite–biotite indicates lower ambient temperature of greenschist facies condition.

Various alternative interpretations regarding the relationship between the metamorphism and deformation in the SSZ have also been put forward (Naha, 1965; Mukhopadhyay et al., 1975; Bhattacharyya and Sanayal, 1988). If higher grade metamorphism in the SSZ and in the northern belt (hanging wall of the SSZ) predates (Naha, 1965), or post-dates (Mukhopadhyay et al., 1975) the deformation responsible for the development of grain-shape fabric and CPO in the SSZ, then activation of the prism  $\langle a \rangle$  slip system and the Y-maximum should have been controlled by factors other than temperature, especially when the CPO from the footwall quartzites are also taken into account.

Alternatively, it has been suggested that temperature may not be the only parameter influencing activation of different slip system (Hobbs, 1985). Fluids may have an important effect on the operative slip systems in natural quartzites (Lister and Dornsiepen, 1982; Bouchez et al., 1984; Blumenfeld et al., 1986). Hippertt (1994a,b) has correlated increasing mica enrichment (phylionitisation) and access of water during progressive strain to a change in operative slip system from basal  $\langle a \rangle$  to prism  $\langle a \rangle$ , through dislocation-assisted diffusion. Prism  $\langle c \rangle$  slip is shown to be facilitated at high temperatures and high fluid pressure (Blacic, 1975; Blumenfeld et al., 1986; Garbutt and Teyssier, 1991). However, to our knowledge, independent influence of the fluid activity on the activation of different slip system in quartzite has not so far been sufficiently evaluated.




Fig. 8. Quartz  $c$ -axis fabric diagrams of the specimens from the Chaibasa Formation immediately north of the Singhbhum shear zone.




Fig. 9. Quartz *c*-axis fabric for the recrystallised quartz grains (column a) vis-a-vis the relict quartz grains (column b). Lower hemisphere equal area projections. *N* is total number of *c*-axis measured in each specimen. Contouring after Kamb method with  $E = 3\sigma$ . Lowest contour corresponds to  $(E - 2\sigma)$ . CI is contour interval. Viewing towards east or southeast. The fabric diagram is not contoured when *N* is less than 100 for relict grains.




Fig. 10. *c*-Axis fabric skeletons of the analysed specimens. Viewing towards east or southeast. Note persistent asymmetry of fabric skeletons, particularly among the SSSZ and the JSZ specimens.

### 7.3. Comparable *c*-axis fabric in recrystallised and relict quartz grains

The *c*-axis fabrics measured from relict quartz grains display a pattern and asymmetry comparable to those from their recrystallised counterparts. Experimental observations demonstrate that CPO of dynamically recrystallised quartzites depends on the mechanism of recrystallisation (Gleason et al., 1993). Dominance of grain-boundary-migration recrystallisation driven by local difference in strain-induced dislocation density leads to CPO of dynamically recrystallised quartz grains distinct from that of deformed original grains (regime I of Hirth and Tullis, 1992). On the other hand dynamic recrystallisation by progressive subgrain rotation leads to a CPO in dynamically recrystallised quartz grains mimicking that of the relict grains because of host control on subgrain orientations (regime II of Hirth and Tullis, 1992). In the regime III of Hirth and Tullis (1992), there is again a preponderance of grain boundary migration, but now driven by surface energy reduction through increased grain boundary mobility at elevated temperature. In regime III as there is no preferential orientation for growth, the CPO of dynamically recrystallised quartz grains is similar to that of relict quartz grains (Gleason et al., 1993). The apparent similarity of *c*-axis fabrics of recrystallised and relict quartz grains from the SSZ and its footwall may be interpreted as a case of regime II or regime III recrystallisation. However, one should keep in mind that fabric patterns or kinematic frame-

work of experimentally deformed quartzites (Gleason et al., 1993) is different from that of the SSZ sample.

### 7.4. Fluid activity in the Singhbhum shear zone

In the present sample, the Y-axis maxima are not associated with an increase in the mica content as suggested by Hippertt (1994a). On the other hand, the role of fluids in the SSZ is demonstrated by the presence of stable hydrous phase such as muscovite and common syntectonic quartz veins. Also the ore minerals in the shear zone are interpreted to be of hydrothermal origin (Sarkar, 1984). Asymmetrically boudinaged quartz veins, whose asymmetry can be related to the syntectonic *C'* shears (Passchier and Trouw, 1996) in the SSZ, are quite common in the mylonites of Singhbhum region (Fig. 3b). This intensity of syntectonic veining could be correlated with a high fluid pressure during deformation. Experimental works demonstrate that the Tuttle lamellae formation is enhanced by either high temperature (Lemmleyn and Kliya, 1960) or by the presence of reactive pore fluid (Smith and Evans, 1984).

Microstructures indicating a coupling of progressive subgrain rotation and grain boundary migration recrystallisation mechanisms, specimens having extensive dynamic recrystallisation and the presence of *S*-*C* fabrics from the SSZ sample may be interpreted as a result of deformation in the regime III or in a transitional regime between regimes II and III (Hirth and Tullis, 1992). Microstructural evolution in such a regime could have occurred at elevated temperature or through the influence of fluids (Hirth and Tullis, 1992). The latter may be the case of water weakening (Blacic, 1975; Mainprice et al., 1986; Paterson, 1989) or transient high fluid pressure leading to brittle effects and consequent water penetration and weakening of quartz. Alternatively, the presence of stable hydrous phases, such as muscovite, indicates that the ambient temperature in the SSZ and JSZ was not too high.

Difficulties remain in attempting to explain the Y-maximum fabric from the SSZ and the JSZ in terms of higher ambient temperature of deformation. On the other hand, experiments (Blacic, 1975) and natural examples of Y-maximum fabric supposedly formed under high fluid activity (e.g. Blumenfeld et al., 1986; cf. Mainprice et al., 1986), suggest that the high partial pressure of water prevalent in the SSZ and its footwall may account for mechanical effects which facilitate water penetration and consequent hydrolytic weakening of quartz leading to the prism  $\langle a \rangle$  slip in addition to the basal  $\langle a \rangle$  and rhomb  $\langle a \rangle$  slips under greenschist facies condition.




Fig. 11. Bar chart indicating frequency of measured quartz *c*-axis fabrics with different combinations of maximum types. Different maximum types (or their combinations) are labelled according to the relative positions of point maximum with respect to the X-, Y- and Z-axes of the finite strain ellipsoid (see text). 'NONE' refers to specimens with no noticeable point concentration in the fabric diagram.




Fig. A1. Comparison of quartz *c*-axis fabric from two mutually perpendicular sections for individual specimens. Orientations measured in sections perpendicular to lineation are rotated before presentation. E–W horizontal line represents the trace of foliation with top of foliation on the north side; arrow at the western end represents down plunge direction of stretching/mineral lineation. Viewing is towards east or southeast. Contouring after Kamb method with  $E = 3\sigma$ . Lowest contour is for  $E = 2\sigma$ . CI is contour interval.

### 7.5. Comparison of *c*-axis fabrics from the SSZ with those from its footwall

The dynamically recrystallised quartz *c*-axis fabrics in the SSZ and the Dhanjori Group quartzites (the footwall rocks of the SSZ) display comparable patterns (Fig. 10), although the actual positions of the fabric point-maxima and the angles made by the leading edge, trailing edge and central segment of the fabrics within individual specimens vary with respect to finite strain axes. Therefore, the deformation responsible for the *c*-axis fabric development in the SSZ and in the Dhanjori Group may be kinematically linked.

## 8. Conclusions

Specimens from the Singhbhum shear zone and its footwall display either asymmetric, type I cross-girdle or asymmetric, kinked single-girdle *c*-axis fabrics measured in dynamically recrystallised quartz grains. The *c*-axis fabric and its asymmetry in the relict quartz grain is apparently very similar to that of the recrystallised quartz grain counterparts. The fabrics are characterised by a concentration near the Y-axis of the finite strain ellipsoid, with an equally well, or subordinately developed, concentration at a small angle to the Z-axis and at an intermediate orientation. The asymmetry of the fabric indicates a general southward thrust sense of movement of the northern block on both the northerly dipping SSZ and subsidiary shear zones such as the JSZ. The formation of quartz CPO with Y-maximum and simultaneous concentration near Z or at an intermediate orientation under greenschist facies condition of the Singhbhum shear zone and its footwall is attributed to influence of higher fluid activity in these large shear zones.

## Acknowledgements

We thank two anonymous reviewers for their critical comments and suggestions on an earlier version, which helped us immensely in improving the paper. Financial support for this work was provided by the Indian Statistical Institute, Calcutta, India.

## Appendix A. Homogeneity of the quartz *c*-axis fabric in the hand specimen scale: A visual check

In order to check the homogeneity of quartz *c*-axis orientations at the hand specimen scale, two thin sections of different orientations were prepared from each specimen—one cut parallel to the mineral elongation lineation and perpendicular to the foliation and the

other perpendicular to both foliation and mineral elongation lineation. *C*-axis orientations of 200 dynamically recrystallised quartz grains were measured in each thin section using a Federov stage fixed to an optical microscope. The measured *c*-axes were then plotted on a lower hemisphere equal area projection and contoured. Necessary rotations were applied so that prepared fabric diagrams refer to the same orientation framework, i.e. with the E–W axis of the fabric diagram as the trace of foliation, the W (left) as the mineral elongation lineation direction, and the north (top) side of the E–W line as the top of the foliation. The viewing direction is towards SE or E. Visual comparison of such fabric diagrams for 12 specimens (a pair for each) show that they are comparable (Fig. A1) and the assumption of homogeneity in quartz *c*-axis fabric at the hand specimen scale could be accepted by the visual comparison method.

## References

- Banerji, A.K., 1959. Cross-folding and thrust tectonics from the Singhbhum shear zone. *Quarterly Journal of the Geological Mining and Metallurgical Society of India* 31, 59–60.
- Bell, T., Etheridge, M.A., 1976. The deformation and recrystallisation of quartz in a mylonite zone, Central Australia. *Tectonophysics* 32, 238–267.
- Bhattacharyya, D.S., Sanayal, P., 1988. The Singhbhum orogen—its structure and stratigraphy. *Memoir of the Geological Society of India* 8, 5–11.
- Blacic, J.D., 1975. Plastic deformation mechanisms in quartz: The effect of water. *Tectonophysics* 27, 271–294.
- Blumenfeld, P., Mainprice, D., Bouchez, J.L., 1986. *C*-slip in quartz from subsolidus deformed granite. *Tectonophysics* 127, 97–115.
- Bose, M.K., 1994. Sedimentation pattern and tectonic evolution of Proterozoic Singhbhum Basin in the Eastern Indian shield. *Tectonophysics* 231, 325–346.
- Bose, M.K., Chakraborty, M.K., 1981. Fossil marginal basin from the Indian shield: A model for the geology of the Singhbhum Precambrian belt, eastern India. *Geologische Rundschau* 70, 504–518.
- Bouchez, J.L., 1977. Plastic deformation of quartzites at low temperature in areas of natural strain gradient. *Tectonophysics* 39, 25–50.
- Bouchez, J.L., Lister, G.S., Nicolas, A., 1983. Fabric asymmetry and shear sense in movement zones. *Geologische Rundschau* 72, 401–419.
- Bouchez, J.L., Mainprice, D.H., Trepied, L., Doukhan, J.C., 1984. Secondary lineation in a high-*T* quartzite (Galicia, Spain): an explanation for an abnormal fabric. *Journal of Structural Geology* 6, 159–165.
- Bunge, H.J., Wenk, H.R., 1977. Three dimensional texture analysis of quartzite (trigonal crystal and triclinic specimen symmetry). *Tectonophysics* 40, 257–285.
- Culshaw, N., 1987. Microstructures, *c*-axis pattern, microstrain and kinematics of some *S*–*C* mylonites in Grenville gneiss. *Journal of Structural Geology* 9, 299–311.
- Davis, G.H., Gardulski, A.F., Lister, G.S., 1987. Shear origin of quartzite mylonite and mylonitic pegmatite in the Coyote Mountain metamorphic core complex, Arizona. *Journal of Structural Geology* 9, 289–297.

- Dunn, J.A., Dey, A.K., 1942. The geology and petrology of eastern Singhbhum and surrounding areas. *Memoir of the Geological Survey of India* 69, 281–456.
- Duebendorfer, E.M., Housten, R.S., 1987. Proterozoic accretionary tectonics at the southern margin of the Archean Wyoming craton. *Geological Society of America Bulletin* 98, 554–568.
- Gall, G., 1964. Precambrian flysch and molasse tectonics and sedimentation around Rakha mines and Jaikon in Singhbhum Dt. Bihar India. XXII International Geological Congress Report 4, 331–356.
- Garbutt, J.M., Teyssier, C., 1991. Prism  $\langle c \rangle$  slip in the quartzites of the Okhurst mylonite belt, California. *Journal of Structural Geology* 13, 657–666.
- Ghosh, S.K., Sengupta, S., 1987a. Structural history of the Singhbhum shear zone in relation to the northern belt. In: Saha, A.K. (Ed.), *Geological Evolution of Peninsular India—Petrological and Structural Aspects, Recent Researches in Geology*, 13. Hindustan Publishing Company, New Delhi, pp. 31–44.
- Ghosh, S.K., Sengupta, S., 1987b. Progressive development of structures in a ductile shear zone. *Journal of Structural Geology* 9, 277–287.
- Geason, G.C., Tullis, J., Heidelbach, H.F., 1993. The role of dynamic recrystallisation in the development of lattice preferred orientation in experimentally deformed quartz aggregates. *Journal of Structural Geology* 15, 1145–1168.
- Gupta, A.K., Basu, A., Ghosh, P.K., 1980. The Proterozoic ultramafic and mafic lavas and tuffs of the Dalma greenstone belt, Singhbhum, eastern India. *Canadian Journal of Earth Science* 17, 210–231.
- Hippert, J.F., 1994a. Microstructures and  $c$ -axis fabrics indicative of quartz dissolution in sheared quartzites and phyllonites. *Tectonophysics* 229, 141–163.
- Hippert, J.F., 1994b. Direct observation of porosity in quartzite and phyllonite. *Neues Jahrbuch für Mineralogie* 166, 239–259.
- Hirth, G., Tullis, J., 1992. Dislocation creep regimes in quartz aggregates. *Journal of Structural Geology* 14, 145–159.
- Hobbs, B.E. 1985. The geological significance of microfabric analysis. In: Wenk, H.R. (Ed.), *Preferred Orientation of Metals and Rocks: An Introduction to Modern Texture Analysis*. Academic Press, Orlando, pp. 463–484.
- Jessell, M.W., 1987. Grain-boundary migration microstructures in a naturally deformed quartzite. *Journal of Structural Geology* 9, 1007–1014.
- Jessell, M.W., 1988a. Simulation of fabric development in recrystallising aggregates—I. Description of the model. *Journal of Structural Geology* 10, 771–778.
- Jessell, M.W., 1988b. Simulation of fabric development in recrystallising aggregates—II. Example model runs. *Journal of Structural Geology* 10, 779–794.
- Jessell, M.W., Lister, G.S., 1990. A simulation of temperature dependence of quartz fabric. In: Knipe, R.J., Rutter, E.H. (Eds.), *Deformation Mechanism, Rheology and Tectonics*, 54, pp. 353–362 Geological Society Special Publication.
- Jiang, D., White, J.C., 1995. Kinematics of flow and the interpretation of geological structures, with special reference to shear zones. *Journal of Structural Geology* 17, 1249–1265.
- Joy, S., 1996. Kinematics of deformation and quartz  $c$ -axis fabric in a part of Singhbhum Shearzone and its footwall, Singhbhum district, Bihar. Ph.D. Thesis, University of Calcutta, Calcutta.
- Krischner, D., Teyssier, C., 1991. Quartz  $c$ -axis fabric difference between porphyroblast and recrystallized grains. *Journal of Structural Geology* 13, 105–109.
- Kutty, T.S., Joy, S., 1997. Sternet—a computer program for stereographic projection: with a new algorithm for contouring. *Journal of the Geological Society of India* 50, 649–653.
- Law, R.D., 1987. Heterogeneous deformation and quartz crystallographic fabric transitions: natural examples from the Stack of Glencoul, northern Assynt. *Journal of Structural Geology* 9, 819–833.
- Law, R.D., Casey, M., Knipe, R.J., 1986. Kinematic and tectonic significance of microstructures and crystallographic fabric within quartz mylonite from the Assynt and Eriboll regions of the Moine thrust zone NW Scotland. *Transactions of the Royal Society of Edinburgh (Earth Science)* 77, 99–126.
- Law, R.D., Knipe, R.J., Dayan, H., 1984. Strain path partitioning within thrust sheets: microstructural and petrofabric evidence from the Moine thrust zone at Loch Eriboll, North-West Scotland. *Journal of Structural Geology* 6, 477–497.
- Lemleyn, G.G., Kliya, M.O., 1960. Distinctive features of the healing of a crack in a crystal under conditions of declining temperature. *International Geology Review* 2, 125–128.
- Lister, G.S., 1977. Cross-girdle  $c$ -axis fabric in quartzites plastically deformed by plane strain and progressive simple shear. *Tectonophysics* 1, 51–54.
- Lister, G.S., Dornsiepen, U.F., 1982. Fabric transition in the Saxony granulite terrain. *Journal of Structural Geology* 1, 283–297.
- Lister, G.S., Hobbs, B.E., 1980. The simulation of fabric development during plastic deformation and its application to quartzite: the influence of deformation history. *Journal of Structural Geology* 2, 355–370.
- Lister, G.S., Price, G.P., 1978. Fabric development in quartz-feldspar mylonite. *Tectonophysics* 49, 37–78.
- Lister, G.S., Snoke, A.W., 1984.  $S$ - $C$  mylonite. *Journal of Structural Geology* 6, 617–638.
- Lister, G.S., Williams, P.F., 1979. Fabric development in shear zone: theoretical controls and observed phenomena. *Journal of Structural Geology* 1, 283–297.
- Lister, G.S., Williams, P.F., 1983. The partitioning of deformation in flowing rock masses. *Tectonophysics* 45, 107–158.
- Mainprice, D., Bouchez, J.L., Blumenfeld, P., Tubia, J.M., 1986. Dominant  $c$ -slip in naturally deformed quartz: implications for dramatic plastic softening at high temperature. *Geology* 14, 819–822.
- Marjoribanks, R.W., 1976. The relation between microfabric and strain in a progressively deformed quartzite sequence from central Australia. *Tectonophysics* 32, 269–293.
- Mukhopadhyay, D., 1976. Precambrian stratigraphy of Singhbhum—the problems and a prospect. *Indian Journal of Earth Science* 3, 208–219.
- Mukhopadhyay, D., Ghosh, A.K., Bhattacharyya, S., 1975. A re-assessment of the structure of the Singhbhum shear zone. *Bulletin of the Geological Mining and Metallurgical Society of India* 48, 49–67.
- Naha, K., 1965. Metamorphism in relation to stratigraphy, structure and movements in parts of east Singhbhum, Eastern India. *Quarterly Journal of the Geological Mining and Metallurgical Society of India* 37, 41–88.
- Nicholas, A., Poirier, J.P., 1976. *Crystalline Plasticity and Solid State Flow in Metamorphic Rocks*. John Wiley and Sons, London.
- Passchier, C.W., 1983. The reliability of asymmetric  $c$ -axis fabrics of quartz to determine sense of vorticity. *Tectonophysics* 99, T9–T18.
- Passchier, C.W., Trouw, R.A.J., 1996. *Microtectonics*. Springer-Verlag, Berlin.
- Paterson, M.S. 1989. The interaction of water with quartz and its influence in dislocation flow—an overview. In: Karato, S., Tourimi, M. (Eds.), *Rheology of Solids and of the Earth*. University Press, New York, pp. 171–194.
- Ralser, S., Hobbs, B.E., Ord, A., 1991. Experimental deformation of a quartz mylonite. *Journal of Structural Geology* 13, 837–850.
- Saha, D., Joy, S., 1995. Dynamic recrystallisation of quartz in the



- Singhbhum shear zone and Dhanjori quartzites, Eastern Singhbhum, Bihar. Abstract, 82nd Indian Science Congress.
- Saha, A.K., Ray, S.L., 1984. The structural and geochemical evolution of the Singhbhum granite batholithic complex, India. *Tectonophysics* 105, 163–176.
- Sarkar, A.N., 1982a. Precambrian tectonic evolution of eastern India; A model for converging micro plates. *Tectonophysics* 86, 363–397.
- Sarkar, A.N., Bhattacharyya, D.S., 1978. Deformation and metamorphism in the western part of the Singhbhum shear belt around Lapsa Buru and Kharsawan, Bihar. *Journal of the Geological Society of India* 19, 310–320.
- Sarkar, S.C. 1966. Structure and control of ore mineralisation in the Moinajharia–Mosaboni–Surda section of the Singhbhum copper belt, Bihar. In: Deb, S. (Ed.), *Contributions to the Geology of Singhbhum*. Jadavpur University, Calcutta, pp. 91–101.
- Sarkar, S.C., 1982b. Uranium (–nickel–cobalt–molybdenum) mineralisation along the Singhbhum copper belt, India and problem of ore genesis. *Mineralium Deposita* 17, 257–278.
- Sarkar, S.C., 1984. *Geology and Ore Mineralisation of the Singhbhum Copper–Uranium Belt, Eastern India*. Jadavpur University, Calcutta.
- Sarkar, S.N., Saha, A.K., 1977. The present status of the Precambrian stratigraphy, tectonics and geochronology of Singhbhum–Keonjhar–Mayurbhanj region, Eastern India. *Indian Journal of Earth Science, S. Ray Volume*, 37–65.
- Schmid, S.M., Casey, M., 1986. Complete fabric analysis of some commonly observed quartz *c*-axis patterns. Mineral and rock deformation: laboratory studies 36, pp. 263–286, *The Paterson Volume*, American Geophysical Union, Geophysical Monograph.
- Simpson, C., Schmid, S.M., 1983. An evaluation of criteria to deduce the sense of movement in sheared rocks. *Geological Society of America Bulletin* 94, 1281–1288.
- Smith, D.L., Evans, B., 1984. Diffusional crack healing in quartz. *Journal of Geophysical Research* 89, 4125–4136.
- Tullis, J.A., 1977. Preferred orientation of quartz produced by slip during plane strain. *Tectonophysics* 39, 87–102.
- Tullis, J.A., Christie, J.M., Griggs, D.T., 1973. Microstructure and preferred orientation of experimentally deformed quartzites. *Geological Society of America Bulletin* 84, 297–314.
- Tuttle, O.F., 1949. Structural petrology of planes of liquid inclusions. *Journal of Geology* 57, 331–356.
- Virnave, S.N., Mukhopadhyay, T.K., Krishnabhadri, S.N.R., 1994. On some aspects of the stratigraphy, depositional environment, and its bearing on uranium mineralisation in parts of the Singhbhum shear zone, Bihar. *Journal of the Geological Society of India* 43, 557–571.
- Vissers, R.L.M., 1993. Quartz *c*-axis fabrics in deformed conglomerates: some support for a skeletal approach to fabric analysis. *Journal of Structural Geology* 15, 1055–1060.
- Wenk, H.R., Canova, G., Molinari, A., Cocks, U.F., 1989. Viscoplastic modelling of texture development in quartzite. *Journal of Geophysical Research* 94, 17895–17906.
- White, S., 1976. The effect of strain on the microstructures, fabrics and deformation mechanisms in quartzite. *Philosophical Transactions Royal Society of London A283*, 69–86.
- White, S., Burrows, S.E., Carreras, J., Shaw, N.D., Humphreys, F.J., 1980. On mylonites in ductile shear zones. *Journal of Structural Geology* 2, 175–187.