

Phase Distribution of Kerr Vectors in a Deformed Hilbert Space

P. K. Das¹

Received August 11, 1998

In this paper we discuss Kerr vectors and their phase distribution in a deformed Hilbert space. We also discuss coherent phase vectors in this space.

1. INTRODUCTION

We consider the set

$$H_q = \{f: f(z) = \sum a_n z^n \text{ where } \sum [n]! |a_n|^2 < \infty\}$$

where $[n] = (1 - q^n)/(1 - q)$, $0 < q < 1$.

For $f, g \in H_q$, $f(z) = \sum_{n=0}^{\infty} a_n z^n$, $g(z) = \sum_{n=0}^{\infty} b_n z^n$ we define addition and scalar multiplication as follows:

$$f(z) + g(z) = \sum_{n=0}^{\infty} (a_n + b_n) z^n \quad (1)$$

and

$$\lambda \circ f(z) = \sum_{n=0}^{\infty} \lambda a_n z^n \quad (2)$$

It is easily seen that H_q forms a vector space with respect to usual pointwise scalar multiplication and pointwise addition by (1) and (2). We observe that $e_q(z) = \sum_{n=0}^{\infty} (z^n/[n]!)$ belongs to H_q .

¹Physics and Applied Mathematics Unit, Indian Statistical Institute, Calcutta 700035, India; e-mail: daspk@isical.ac.in.

Now we define the inner product of two functions $f(z) = \sum a_n z^n$ and $g(z) = \sum b_n z^n$ belonging to H_q as

$$(f, g) = \sum [n]! \bar{a}_n b_n \quad (3)$$

The corresponding norm is given by

$$\|f\|^2 = (f, f) = \sum [n]! |a_n|^2 < \infty$$

With this norm derived from the inner product it can be shown that H_q is a complete normed space. Hence H_q forms a Hilbert space.

In a recent paper⁽¹⁾ we proved that the set $\{z^n / \sqrt{[n]!}, n = 0, 1, 2, 3, \dots\}$ forms a complete orthonormal set. If we consider the following actions on H_q :

$$\begin{aligned} T f_n &= \sqrt{[n]} f_{n-1} \\ T^* f_n &= \sqrt{[n+1]} f_{n+1} \end{aligned} \quad (4)$$

where T is the backward shift and its adjoint T^* is the forward shift operator on H_q and $f_n(z) = z^n / \sqrt{[n]!}$, then we have shown⁽¹⁾ that the solution of the following eigenvalue equation

$$T f_\alpha = \alpha f_\alpha \quad (5)$$

is given by

$$f_\alpha = e_q(|\alpha|^2)^{-1/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{[n]!}} f_n \quad (6)$$

We call f_α a *coherent vector* in H_q .

This paper is divided into four sections. In Section 1 we have given an introduction stating coherent vectors in H_q . In section 2 we introduce Kerr vectors in H_q . Section 3 deals with phase distributions in H_q and in Section 4 we discuss coherent phase vectors in this space.

2. GENERATION OF KERR VECTORS

The Kerr vectors in H_q are defined by

$$\phi_\alpha^K = e_q^{(i/2) \gamma [N(N-1)]} f_\alpha \quad (7)$$

where $f_\alpha \in H_q$ is a coherent vector given by (6), γ is a constant, and $N = T^*T$, with T the backward shift (4).

Now,

$$\begin{aligned}
 \phi_{\alpha}^K &= e_q^{(i/2)\gamma N(N-1)} f_{\alpha} \\
 &= e_q^{(i/2)\gamma N(N-1)} e_q(|\alpha|^2)^{-1/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n}!} f_n \\
 &= e_q(|\alpha|^2)^{-1/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n}!} e_q^{(i/2)\gamma[n]([n]-1)} f_n \\
 &= \sum_{n=0}^{\infty} \left[e_q(|\alpha|^2)^{-1/2} \frac{\alpha^n}{\sqrt{n}!} e_q^{(i/2)\gamma[n]([n]-1)} \right] f_n \\
 &= \sum_{n=0}^{\infty} q_n f_n
 \end{aligned} \tag{8}$$

where

$$q_n = e_q(|\alpha|^2)^{-1/2} \frac{\alpha^n}{\sqrt{n}!} e_q^{(i/2)\gamma[n]([n]-1)} \tag{9}$$

The *quasiprobability distribution*, known as the *Q function*, for Kerr vectors (7) is then defined by

$$\begin{aligned}
 Q(\beta) &= |(f_{\beta}, \phi_{\alpha}^K)|^2 \\
 &= |e_q(|\alpha|^2)^{-1/2} \sum_{n=0}^{\infty} \frac{\bar{(\beta)}^n}{\sqrt{n}!} q_n|^2
 \end{aligned} \tag{10}$$

where q_n is given by (9).

3. PHASE DISTRIBUTION

To obtain the phase distribution we consider first the *phase operator* $P = (q^n + T^*T)^{-1/2}T$ and try to find the solution of the following eigenvalue equation:

$$Pf_{\beta} = \beta f_{\beta} \tag{11}$$

$$f_{\beta}(z) = \sum_{n=0}^{\infty} a_n z^n = \sum_{n=0}^{\infty} a_n \sqrt{n}! f_n(z) \tag{12}$$

$$\text{or, } f_{\beta} = \sum_{n=0}^{\infty} a_n \sqrt{n}! f_n$$

$$Pf_{\beta} = \sum_{n=0}^{\infty} a_n \sqrt{n}! (q^n + T^*T)^{-1/2} T f_n$$

$$\begin{aligned}
&= \sum_{n=1}^{\infty} a_n \sqrt{n!} (q^n + T^*T)^{-1/2} \sqrt{n!} f_{n-1} \\
&= \sum_{n=1}^{\infty} a_n \sqrt{n!} \sqrt{n!} (q^n + [n-1])^{-1/2} f_{n-1} \\
&= \sum_{n=0}^{\infty} a_{n+1} \sqrt{n+1!} \sqrt{n+1!} (q^{n+1} + [n])^{-1/2} f_n \quad (13)
\end{aligned}$$

$$\beta f_{\beta} = \beta \sum_{n=0}^{\infty} a_n \sqrt{n!} f_n \quad (14)$$

From (11)–(14) we observe that a_n satisfies the following difference equation:

$$a_{n+1} \sqrt{n+1!} \sqrt{n+1!} (q^{n+1} + [n])^{-1/2} = \beta a_n \sqrt{n!} \quad (15)$$

That is,

$$a_{n+1} = \frac{\beta a_n (q^{n+1} + [n])^{1/2}}{[n+1]} \quad (16)$$

Hence,

$$\begin{aligned}
a_1 &= \frac{\beta(q + [0])^{1/2} a_0}{[1]} \\
a_2 &= \frac{\beta a_1 (q^2 + [1])^{1/2}}{[2]} = \frac{\beta^2 a_0 \sqrt{(q + [0])(q^2 + [1])}}{[2]!} \\
a_3 &= \frac{\beta a_2 (q^3 + [2])^{1/2}}{[3]} = \frac{\beta^3 a_0 \sqrt{(q + [0])(q^2 + [1])(q^3 + [2])}}{[3]!}
\end{aligned}$$

and so on. Thus,

$$a_n = \frac{\beta^n a_0 \sqrt{(q + [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}}{[n]!}$$

Hence,

$$\begin{aligned}
f_{\beta} &= \sum_{n=0}^{\infty} a_n \sqrt{n!} f_n \\
&= a_0 \sum_{n=0}^{\infty} \beta^n \frac{\sqrt{(q + [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}}{[n]!} f_n
\end{aligned}$$

where $\beta = |\beta|e^{i\theta}$ is a complex number. These vectors are normalizable in a strict sense only for $|\beta| < 1$.

Now, if we take $a_0 = 1$ and $|\beta| = 1$, we have

$$f_\beta = \sum_{n=0}^{\infty} e^{in\theta} \sqrt{\frac{(q+ [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!}} f_n \quad (17)$$

Henceforth, we shall denote this vector as

$$f_\theta = \sum_{n=0}^{\infty} e^{in\theta} \sqrt{\frac{(q+ [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!}} f_n \quad (18)$$

$0 \leq \theta \leq 2\pi$, and we call f_θ a *phase vector* in H_q .

The phase vectors f_θ are neither normalizable nor orthogonal. The completeness relation

$$I = \frac{1}{2\pi} \int_0^{2\pi} d\mu(\theta) |f_\theta\rangle\langle f_\theta| \quad (19)$$

where

$$d\mu(\theta) = \frac{[n]!}{(q+ [0])(q^2 + [1]) \dots (q^n + [n-1])} d\theta \quad (20)$$

may be proved as follows:

Define the operator

$$|f_\theta\rangle\langle f_\theta|: H_q \rightarrow H_q \quad (21)$$

by

$$|f_\theta\rangle\langle f_\theta|f = (f_\theta, f)f_\theta \quad (22)$$

with

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

Now,

$$\begin{aligned} & (f_\theta, f) \\ &= \sum_{n=0}^{\infty} [n]! \frac{e^{-in\theta}}{\sqrt{[n]!}} \sqrt{\frac{(q+ [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!}} a_n \\ &= \sum_{n=0}^{\infty} e^{-in\theta} \sqrt{(q+ [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])} a_n \end{aligned} \quad (23)$$

Then,

$(f_\theta, f)f_\theta$

$$= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} a_n e^{i(m-n)\theta} \frac{\sqrt{(q+[0])(q^2+[1]) \dots (q^m+[m-1])}}{[m]!} \quad (24)$$

$$\times \sqrt{(q+[0])(q^2+[1]) \dots (q^n+[n-1])} f_m$$

Using

$$\int_0^{2\pi} d\theta e^{i(m-n)\theta} = 2\pi\delta_{nm} \quad (25)$$

we have

$$\frac{1}{2\pi} \int_0^{2\pi} d\mu(\theta) |f_\theta\rangle\langle f_\theta| f$$

$$= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} a_n f_m \frac{\sqrt{(q+[0])(q^2+[1]) \dots (q^m+[m-1])}}{[m]!}$$

$$\times \sqrt{(q+[0])(q^2+[1]) \dots (q^n+[n-1])} \frac{1}{2\pi} \int_0^{2\pi} e^{i(m-n)\theta} d\theta$$

$$\times \frac{[n]!}{(q+[0])(q^2+[1]) \dots (q^n+[n-1])}$$

$$= \sum_{n=0}^{\infty} a_n f_n \frac{(q+[0])(q^2+[1]) \dots (q^n+[n-1])}{\sqrt{[n]!}}$$

$$\times \frac{[n]!}{(q+[0])(q^2+[1]) \dots (q^n+[n-1])}$$

$$= \sum_{n=0}^{\infty} \sqrt{[n]!} a_n f_n$$

$$= f \quad (26)$$

Thus, (19) follows.

The *phase distribution* over the window $0 \leq \theta \leq 2\pi$ for any vector f is then defined by

$$P(\theta) = \frac{1}{2\pi} |(f_\theta, f)|^2 \quad (27)$$

For the Kerr vector, $P(\theta)$ is given by

$$\begin{aligned}
 P(\theta) &= \frac{1}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_n \bar{q}_m e^{i(m-n)\theta} \sqrt{(q+[0])(q^2+[1])(q^3+[2]) \dots (q^n+[n-1])} \\
 &\quad \times \sqrt{(q+[0])(q^2+[1])(q^3+[2]) \dots (q^m+[m-1])}
 \end{aligned} \tag{28}$$

where q_n is given by (9).

4. COHERENT PHASE VECTORS

The coherent phase vectors $\{f_\beta\}$ are the unit-length eigenvectors of the operator P of (11) whose eigenvalues β are complex numbers, namely,

$$P f_\beta = \beta f_\beta$$

From the previous section we see that

$$\begin{aligned}
 f_\beta &= \sum_{n=0}^{\infty} a_n \sqrt{n!} f_n \\
 &= a_0 \sum_{n=0}^{\infty} \beta^n \sqrt{\frac{(q+[0])(q^2+[1])(q^3+[2]) \dots (q^n+[n-1])}{[n!]}} f_n
 \end{aligned} \tag{29}$$

Here, $|\beta| < 1$ is necessary in order to ensure normalization, for,

$$\|f_\beta\|^2 = |a_0|^2 \sum_{n=0}^{\infty} |\beta|^{2n} \frac{(q+[0])(q^2+[1])(q^3+[2]) \dots (q^n+[n-1])}{[n!]} \tag{30}$$

The above series (30) is convergent for $|\beta| < 1$. Thus, when f_β is normalized we have $|a_0| = \Phi(|\beta|^2)^{-1/2}$, where

$$\Phi(|\beta|^2) = \sum_{n=0}^{\infty} |\beta|^{2n} \frac{(q+[0])(q^2+[1])(q^3+[2]) \dots (q^n+[n-1])}{[n!]} \tag{31}$$

Thus, f_{β} (29) takes the form

$$f_{\beta} = \Phi(|\beta|^2)^{-1/2} \sum_{n=0}^{\infty} \beta^n \frac{(q + [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!} f_n \quad (32)$$

where $|\beta| < 1$.

It is proper to call these vectors *coherent phase vectors* because they are eigenvectors of the *phase operator* P and they are *coherent* with respect to P .

We have that the coherent phase vectors are normalized and nonorthogonal,

$$(f_{\alpha}, f_{\beta}) = \frac{\Phi(\bar{\alpha}\beta)}{\sqrt{\Phi(|\alpha|^2)\Phi(|\beta|^2)}} \quad (33)$$

If we ignore the normalization factor $\Phi(|\beta|^2)^{-1/2}$, then we get

$$f_{\beta} \rightarrow \sum_{n=0}^{\infty} e^{in\theta} \frac{(q + [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!} f_n = f_{\theta} \quad (34)$$

for $\beta = |\beta|e^{i\theta}$ with $|\beta| \rightarrow 1$.

The *phase representation* of the coherent phase vector $f(\beta)$ is

$$(f_{\theta}, f_{\beta}) = \Phi(|\beta|^2)^{-1/2} \sum_{n=0}^{\infty} \beta^n e^{-in\theta} \frac{(q + [0])(q^2 + [1])(q^3 + [2]) \dots (q^n + [n-1])}{[n]!} = \frac{\Phi(e^{-i\theta}\beta)}{\sqrt{\Phi(|\beta|^2)}} \quad (35)$$

Now, from (35) we have the *phase distribution* over the window $0 \leq \theta \leq 2\pi$ for the *coherent phase vector* f_{β} as

$$\begin{aligned} P(\theta) &= \frac{1}{2\pi} |(f_{\theta}, f_{\beta})|^2 \\ &= \frac{1}{2\pi} \frac{|\Phi(e^{-i\theta}\beta)|^2}{\Phi(|\beta|^2)} \end{aligned} \quad (36)$$

REFERENCES

1. P. K. Das, *Eigenvectors of backwardshift on a deformed Hilbert space*, *Int. J. Theor. Phys.*, **37**, (1998).

2. R. W. Gray and C. A. Nelson, *A completeness relation for the q -analogue coherent states by q -integration*, *J. Phys. A Math. Gen.* **23**, L945–L950 (1990).
3. P. Carruthers and M. M. Nieto, *Phase and angle variables in quantum mechanics*, *Rev. Mod. Phys.* **40**, 411–440 (1968).
4. J. H. Shapiro and S. C. Shepard, *Quantum phase measurement: A system-theory perspective*, *Phys. Rev. A* **43**, 3795–3818 (1991).
5. A. D. Wilson-Gordon, V. Buzek, and P. L. Knight, *Statistical and phase properties of displaced Kerr states*, *Phys. Rev. A* **44**, 7647–7656 (1991).