



## INDIAN STAT

TRANSITIVITY OF PROXIMALITY AND  
NORM ATTAINING FUNCTIONALS

BY

DARAPANENI NARAYANA and T. S. S. R. K. RAO (Bangalore)

**Abstract.** We study the question of when the set of norm attaining functionals on a Banach space is a linear space. We show that this property is preserved by factor reflexive proximal subspaces in  $R(1)$  spaces and generally by taking quotients by proximal subspaces. We show, for  $\mathcal{K}(\ell_2)$  and  $c_0$ -direct sums of families of reflexive spaces, the transitivity of proximality for factor reflexive subspaces. We also investigate the linear structure of the set of norm attaining functionals on hyperplanes of  $c_0$  and show that, for some particular hyperplanes of  $c_0$ , linearity and orthogonal linearity coincide for the set of norm attaining functionals.

**1. Introduction.** We work only with real Banach spaces. For a Banach space  $X$ , we denote by  $B_X$ ,  $S_X$  and  $NA(X)$  the closed unit ball of  $X$ , unit sphere of  $X$  and the set of all norm attaining functionals on  $X$  respectively. For a closed subspace  $Y$  of  $X$  we denote by  $Q_Y$  the canonical quotient map of  $X$  to  $X/Y$ . We are interested in Banach spaces for which  $NA(X)$  is a linear space. It is known that this is intimately related to the question of transitivity of proximality ([4], [7]). We recall that  $Y$  is said to be a *proximal subspace* of  $X$  if for every  $x \in X$  there exists  $y \in Y$  such that  $\|x - y\| = d(x, Y)$ , we then write  $Y \overset{P}{\subseteq} X$ .

In [9] W. Pollul raised the following question on *transitivity of proximality*.

- (A) Which Banach spaces  $X$  have the following property: For any closed subspaces  $Y$  and  $Z$  of  $X$  with  $Y \subseteq Z$ , if  $\dim(X/Z) = \dim(Z/Y) = 1$  and  $Y \overset{P}{\subseteq} Z$ ,  $Z \overset{P}{\subseteq} X$ , then  $Y \overset{P}{\subseteq} X$ ?

In [7] V. Indumathi asked a more general question.

- (B) Which Banach spaces  $X$  have the following property: For any closed subspaces  $Y$  and  $Z$  of  $X$  with  $Y \subseteq Z$ , if  $\dim(X/Y) = n < \infty$  and  $Y \overset{P}{\subseteq} Z$ ,  $Z \overset{P}{\subseteq} X$ , then  $Y \overset{P}{\subseteq} X$ ?

2000 *Mathematics Subject Classification*: 41A65, 41A50, 46B20.

*Key words and phrases*:  $c_0$ , norm attaining functional, orthogonal linearity,  $P$  space,  $R(1)$  space, transitivity of proximality.

Following [7] we call a Banach space  $X$  with property described in (B) a  $P(n)$  space, and we call  $X$  a  $P$  space if it is a  $P(n)$  space for every  $n \geq 2$ . Examples of  $P$  spaces are  $c_0$  and  $\mathcal{K}(\ell_2)$  (the space of compact operators on  $\ell_2$ ). Also any finite-codimensional proximal subspace of a  $P$  space is a  $P$  space ([8]).

A Banach space  $X$  is said to be an  $R(1)$  space if every closed subspace  $Y$  of  $X$  of finite codimension with  $Y^\perp \subseteq \text{NA}(X)$  is proximal in  $X$ . Examples of  $R(1)$  spaces include  $c_0$ , all closed subspaces of  $c_0$ , reflexive spaces and  $\mathcal{K}(\ell_2)$  (see [3] for  $c_0$  and [4] for  $\mathcal{K}(\ell_2)$ ).

To describe the connection between  $R(1)$  and  $P$  spaces we need to recall the concept of orthogonal linearity from [7].

Let  $f, g \in X^*$ . Then  $f$  is said to be *strongly orthogonal* to  $g$  if the supremum of  $f$  on the unit ball of  $X$  is attained at some point of the unit ball of  $\ker g$ . A subset  $F \subset X^*$  is said to be *orthogonally linear* if  $f, g \in F$  and  $f$  strongly orthogonal to  $g$  implies that  $\text{span}\{f, g\} \subseteq F$ . Recall that [7, Question 1] it is not known if there is a space  $X$  for which  $\text{NA}(X)$  is orthogonally linear but not linear. We answer this question in the case of hyperplanes of  $c_0$ .

It was proved in [7] that  $X$  is an  $R(1)$  space and  $\text{NA}(X)$  is orthogonally linear if and only if  $X$  is a  $P$  space. Recently these properties were studied in [8] for direct sums of Banach spaces.

So far we have assumed that the subspaces are of finite codimension. We now consider subspaces with reflexive quotient, called factor reflexive spaces. Thus a closed subspace  $Y$  of a Banach space  $X$  is *factor reflexive* if  $X/Y$  is reflexive. Analogous to the above definitions, we call a Banach space  $X$  an  $\widetilde{R}(1)$  space if for every factor reflexive subspace  $Y$  the condition  $Y^\perp \subseteq \text{NA}(X)$  implies that  $Y$  is proximal in  $X$ . Since any reflexive quotient of  $c_0$  is finite-dimensional,  $c_0$  is an  $\widetilde{R}(1)$  as well as  $\widetilde{R}(1)$  space.

We can now ask the following generalized version of questions (A) and (B).

- (C) Which Banach spaces  $X$  have the following property: For any factor reflexive closed subspaces  $Y$  and  $Z$  of  $X$  with  $Y \subseteq Z$ , if  $Y \overset{P}{\subseteq} Z$  and  $Z \overset{P}{\subseteq} X$  then  $Y \overset{P}{\subseteq} X$ ?

A Banach space with the property in (C) will be called a  $\widetilde{P}$  space. Clearly any reflexive space and the space  $c_0$  are examples of  $\widetilde{P}$  spaces. Also any factor reflexive proximal subspace of a  $\widetilde{P}$  space is again a  $\widetilde{P}$  space.

One of the aims of the present article is to contribute to the study of  $\widetilde{R}(1)$  and  $\widetilde{P}$  spaces. We now briefly describe the content of the article section-wise.

The second section contains investigations on the vector space structure of the norm attaining functionals on a Banach space  $X$ . In particular we study this for a factor reflexive proximal subspace  $Y$  of a  $\widetilde{P}$  space and for

the quotient space  $X/Y$  of a  $P$  space  $X$ . We also give some stability results when  $X$  is an  $\widetilde{R}(1)$  space and  $\text{NA}(X)$  is a vector space.

Motivated by Lemma 4.2 of [4] which identifies  $\text{NA}(\mathcal{K}(\ell_2))$  with the set of finite rank operators, in the third section we show that for any closed subspace of  $\text{NA}(\mathcal{K}(\ell_2))$  (by this we *always* mean that these subspaces are Banach spaces) the pre-annihilator is proximal in  $\mathcal{K}(\ell_2)$ . We also show that  $\mathcal{K}(\ell_2)$  and the  $c_0$ -direct sum of any family of reflexive spaces are  $\widetilde{P}$  spaces.

In the fourth section we show that any separable  $\widetilde{R}(1)$  space can be renormed with a Gateaux smooth norm retaining the proximinality properties. In particular we show that if  $X$  is a separable  $\widetilde{R}(1)$  space then there exists an equivalent Gateaux smooth norm on  $X$  such that  $X$  with this new norm is still an  $\widetilde{R}(1)$  space.

In the fifth section we study the vector space structure of norm attaining functionals in hyperplanes of  $c_0$ . We prove that orthogonal linearity and linearity are equivalent for hyperplanes in  $c_0$ , which gives a partial answer to Question 1 of [7].

**Acknowledgements.** We thank Professors G. Godefroy and G. Pisier for the discussions we had with them while working on this paper. We also thank Professor G. Skandalis for his help in proving Proposition 3.4. The second named author's research was partially supported by an Indo-French project, I.F.C.P.A.R. Grant No. 2601-1. We thank the referee for his extensive comments which improved the readability of the paper.

**2. Linearity of  $\text{NA}(Y)$  for a closed subspace  $Y$  of a Banach space  $X$ .** We start by recalling Garkavi's characterization for finite-codimensional proximal subspaces which we use frequently.

**LEMMA 2.1** (Garkavi [10]). *Let  $X$  be a normed linear space and  $Y$  be a closed subspace of finite codimension. Then  $Y$  is proximal in  $X$  if and only if every closed subspace  $Z \supseteq Y$  of  $X$  is proximal in  $X$ .*

**LEMMA 2.2.** *Let  $Y$  be a proximal subspace of  $X$ . Then  $Y$  is factor reflexive in  $X$  if and only if  $Y^\perp \subseteq \text{NA}(X)$ .*

*Proof.* Suppose  $Y$  is factor reflexive in  $X$ . Equivalently,  $(X/Y)^* \simeq Y^\perp$  is reflexive. Thus every  $f \in Y^\perp$  is norm attaining on  $X/Y$ . Since  $Y$  is proximal in  $X$ ,  $f \in \text{NA}(X)$ . Thus  $Y^\perp \subseteq \text{NA}(X)$ . Conversely, suppose  $Y^\perp \subseteq \text{NA}(X)$ . Then every element  $f$  in  $Y^\perp$  attains its norm on  $X/Y$ . By a well known theorem of James we conclude that  $X/Y$  is reflexive. ■

We now prove the following extension of Garkavi's characterization of finite-codimensional proximal subspaces to the factor reflexive case.

We now turn our attention to the linear structure of the set of norm attaining functionals for quotient spaces by proximal subspaces.

**THEOREM 2.10.** *Let  $Y$  be a proximal subspace of  $X$ .*

- (i) *If  $NA(X)$  is orthogonally linear then so is  $NA(X/Y)$ .*
- (ii) *If  $NA(X)$  is linear then so is  $NA(X/Y)$ .*

*Proof.* We only need to prove (i). Suppose  $NA(X)$  is orthogonally linear. Let  $f, g \in NA(X/Y)$  with  $\|f\| = \|g\| = 1$ . Let  $f', g' \in Y^\perp \subset X^*$  be such that  $Q^*(f) = f'$  and  $Q^*(g) = g'$  where  $Q^* : (X/Y)^* = Y^\perp \rightarrow X^*$  is the inclusion map. Then  $\|f'\| = \|g'\| = 1$  and  $f', g' \in NA(X)$ . We also have  $\|f + g\| = \|f' + g'\|$ .

Suppose  $f$  is strongly orthogonal to  $g$ . Let  $z \in S_{\ker g}$  be such that  $f(z) = 1$ . By the proximality of  $Y$  there exists  $y \in Q_Y(z)$  such that  $\|y\| = 1$  and  $f'(y) = 1$ . It is easily seen that  $y \in \ker g'$ , which implies that  $f'$  is strongly orthogonal to  $g'$ . By orthogonal linearity of  $NA(X)$ , we have  $f' + g' \in NA(X)$ . Hence there is  $x_0 \in S_X$  such that  $(f' + g')(x_0) = \|f' + g'\| = \|f + g\| = (f + g)(Q_Y(x_0))$ , which implies that  $f + g \in NA(X/Y)$ . Hence  $NA(X/Y)$  is orthogonally linear. ■

The following lemma is known. For completeness we give an easy proof.

**LEMMA 2.11.** *Let  $X$  be a Banach space and  $Y$  be a closed subspace. Let  $Z$  be a closed subspace of  $X/Y$ . If  $Q_Y^{-1}(Z)$  is proximal in  $X$ , then  $Z$  is proximal in  $X/Y$ .*

*Proof.* Let  $Q_Y(x_0) \in X/Y$ . Since  $Q_Y^{-1}(Z)$  is proximal, there exists  $z_0 \in Q_Y^{-1}(Z)$  such that  $d(x_0, Q_Y^{-1}(Z)) = \|x_0 - z_0\|$ . Now for any  $z \in Q_Y^{-1}(Z)$  and for any  $n \geq 1$ , we have

$$\begin{aligned} \|Q_Y(x_0 - z)\| &= d(x_0 - z, Y) > \|x_0 - z - y_n\| - 1/n \\ &\geq d(x_0, Q_Y^{-1}(Z)) - 1/n = \|x_0 - z_0\| - 1/n \\ &\geq \|Q_Y(x_0 - z_0)\| - 1/n \end{aligned}$$

where  $y_n \in Y$ . So  $\|Q_Y(x_0 - z)\| \geq \|Q_Y(x_0 - z_0)\|$  for every  $Q_Y(z) \in Z$ , which implies proximality of  $Z$  at  $Q_Y(x_0)$ . Since  $Q_Y(x_0)$  is arbitrary,  $Z$  is proximal in  $X/Y$ . ■

A consequence of the following theorem and the results proved above is that if  $X$  is a  $P$  space and  $Y \subset X$  is reflexive then  $X/Y$  is a  $P$  space.

**THEOREM 2.12.** *Let  $X$  be an  $R(1)$  space and let  $Y$  be a proximal subspace of  $X$ . Then  $X/Y$  is an  $R(1)$  space. Hence if  $X$  is a  $P$  space so is  $X/Y$ .*

*Proof.* Let  $Z$  be a closed subspace of finite codimension  $n$  in  $X/Y$  with  $Z^\perp \subseteq NA(X/Y)$ . Let  $f_1, \dots, f_n \in NA(X/Y)$  be such that  $Z = \bigcap_{i=1}^n \ker f_i$ . Then  $Q_Y^{-1}(Z) = \bigcap_{i=1}^n \ker Q^*(f_i)$  and  $Q^*(f_i) \in Q^*(Z^\perp) \subseteq X^*$ . Since  $Y$  is proximal, we have  $Q^*(Z^\perp) = (Q_Y^{-1}(Z))^\perp \subseteq NA(X)$ . Since  $X$  is an  $R(1)$

space,  $Q_Y^{-1}(Z)$  is proximal in  $X$ . Thus by Lemma 2.11,  $Z$  is proximal in  $X/Y$  and this shows that  $X/Y$  is an  $R(1)$  space. By Theorem 2.10(i), it follows that  $X/Y$  is a  $P$  space if  $X$  is. ■

We do not know an answer to the following version of the "3-space" problem. If  $Y \subset X$  is reflexive and  $X/Y$  is a  $P$  (or  $R(1)$ ) space, is  $X$  a  $P$  space ( $R(1)$  space)?

For a Banach space  $X$ , let  $\mathcal{P}_X = \{Y \stackrel{P}{\subseteq} X : \dim(X/Y) < \infty\}$ . We have the following stability properties.

LEMMA 2.13. *Let  $X$  be an  $R(1)$  space. Then the following statements are equivalent.*

- (i)  $\text{NA}(X)$  is linear.
- (ii)  $\mathcal{P}_X$  is stable under intersection.
- (iii) For any two proximal hyperplanes  $Y_1$  and  $Y_2$  of  $X$ ,  $Y_1 \cap Y_2$  is a proximal subspace of  $X$ .

*Proof.* (i)  $\Rightarrow$  (ii). Assume that  $\text{NA}(X)$  is a vector space. Let  $Y_1$  and  $Y_2$  be two finite-codimensional proximal subspaces of  $X$  with codimensions  $n_1$  and  $n_2$  respectively. Let  $Y_1 = \bigcap_{i=1}^{n_1} \ker f_i$  and  $Y_2 = \bigcap_{j=1}^{n_2} \ker g_j$ , where  $f_1, \dots, f_{n_1}, g_1, \dots, g_{n_2} \in \text{NA}(X)$ . By Garkavi's lemma we have  $Y_1^\perp, Y_2^\perp \subseteq \text{NA}(X)$ . Since  $\text{NA}(X)$  is a vector space and  $Y_1^\perp$  and  $Y_2^\perp$  are finite-dimensional spaces, we have  $\text{span}\{Y_1^\perp, Y_2^\perp\} = \text{span}\{f_1, \dots, f_{n_1}, g_1, \dots, g_{n_2}\} \subseteq \text{NA}(X)$ . Since  $(Y_1 \cap Y_2)^\perp = \text{span}\{Y_1^\perp, Y_2^\perp\}$  and  $X$  is an  $R(1)$  space this implies that  $Y_1 \cap Y_2$  is proximal in  $X$ .

(ii)  $\Rightarrow$  (iii) is trivial.

(iii)  $\Rightarrow$  (i). Suppose  $\text{NA}(X)$  is not linear. Then there exist  $f$  and  $g$  in  $\text{NA}(X)$  such that  $f + g \notin \text{NA}(X)$ . Thus  $\ker f \cap \ker g$  is not proximal in  $X$  (by Garkavi's lemma). ■

PROPOSITION 2.14. *Let  $X$  be an  $R(1)$  space such that  $\text{NA}(X)$  is a linear space. Let  $Y$  be a closed subspace of  $X$ . Then  $\mathcal{P}_Y \subseteq \mathcal{P}_X \cap Y$ . More precisely if  $Z \stackrel{P}{\subseteq} Y$  and  $\dim(Y/Z) = n$ , then there exists a proximal subspace  $Z_0$  in  $X$  of codimension  $n$  such that  $Z = Z_0 \cap Y$ .*

*Proof.* Let  $Z$  be a proximal subspace of  $Y$  of codimension  $n$ . Then by Garkavi's lemma,  $Z^\perp \subseteq \text{NA}(Y) \subseteq Y^*$ . Let  $\{y_i^*\}_{1 \leq i \leq n}$  be a basis of  $Z^\perp$ . Let  $x_i^*$  in  $X^*$  be such that  $x_i^*|_Y = y_i^*$  and  $\|x_i^*\| = \|y_i^*\|$ . This implies that  $x_i^* \in \text{NA}(X)$  for every  $i = 1, \dots, n$ . If  $V = \text{span}\{x_i^* : 1 \leq i \leq n\}$ , then  $V \subseteq \text{NA}(X)$  since  $\text{NA}(X)$  is a vector space. Now  $V^\perp = Z_0 \stackrel{P}{\subseteq} X$  since  $X$  is an  $R(1)$  space. Finally,  $Z_0 \cap Y = Z$ . ■

REMARK 2.15. If  $X$  is an  $R(1)$  space such that  $\text{NA}(X)$  is a vector space and if  $\mathcal{P}_Y = \mathcal{P}_X \cap Y$ , then  $Y$  is also an  $R(1)$  space such that  $\text{NA}(Y)$  is a vector

space. But the converse is not true. For example, let  $X = c_0$  and  $Y = \ker f$  where  $f = (1, 1/3, 1/4, 1/8, \dots) \in \ell_1$ . It will be shown in Proposition 5.4 that  $NA(Y)$  is a vector space. It was shown in [3] that every closed subspace of  $c_0$  is an  $R(1)$  space, which implies that  $\mathcal{P}_Y$  is stable under intersections. Let  $e_1 = (1, 0, 0, \dots) \in \ell_1$ . It is easy to see that  $\ker e_1$  is proximal in  $X$  but  $\ker e_1 \cap Y$  is not in  $\mathcal{P}_Y$ , which implies that  $\mathcal{P}_Y \subsetneq \mathcal{P}_X \cap Y$ .

**3.  $\widetilde{R(1)}$  spaces.** Recall that the rank of an operator  $A : X \rightarrow X$  is the dimension of its image. The following proposition gives a characterization of subspaces of tensor product spaces when the ranks of the elements in the subspace are uniformly bounded.

**PROPOSITION 3.1.** *Let  $E$  and  $F$  be vector spaces and  $E'$  be the algebraic dual of  $E$ . Let  $E' \otimes F = R(E, F)$  be the space of finite rank linear maps from  $E$  to  $F$ . Let  $V$  be a vector subspace of  $R(E, F)$  such that*

$$\sup\{\text{rank}(T) : T \in V\} = N < \infty.$$

*Then there exist  $f_1, \dots, f_N$  in  $F$  and  $e_1^*, \dots, e_N^*$  in  $E'$  such that every  $T$  in  $V$  can be written as*

$$T = \sum_{i=1}^N e_i^* \otimes g_i + \sum_{j=1}^N b_j^* \otimes f_j$$

*for some  $g_1, \dots, g_N \in F$  and  $b_1^*, \dots, b_N^* \in E'$ .*

*Proof.* Let  $T_0$  in  $V$  be such that  $\text{rank}(T_0) = \sup\{\text{rank}(T) : T \in V\} = N < \infty$ . There exists a basis  $\mathcal{B}_1$  of  $E$  and a basis  $\mathcal{B}_2$  of  $F$  such that the matrix of  $T_0$  relative to  $\mathcal{B}_1$  and  $\mathcal{B}_2$  is

$$\left[ \begin{array}{cccc|c} 1 & 0 & \cdots & 0 & \\ 0 & 1 & \cdots & 0 & \\ \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & \cdots & 1 & \\ \hline & & & 0 & 0 \end{array} \right]_{(N \times N)}$$

Let  $V_1 = \{T \in V : x_{pq} = 0 \text{ for all } p \leq N \text{ and } q \leq N\}$  where  $(x_{pq})_{p,q}$  is the matrix of  $T$  with respect to the bases  $\mathcal{B}_1$  and  $\mathcal{B}_2$ . Since  $\dim(V/V_1) < \infty$  and  $V$  consists of finite rank operators, the lemma follows easily from the following claim.

**CLAIM.** *If  $T \in V_1$  and  $i, j \notin \{1, \dots, N\}$ , then  $x_{ij} = 0$ .*

*Proof of the Claim.* Pick  $0 \neq \lambda \in \mathbb{C}$  and let  $S = T_0 + \lambda T$ . We consider the determinant of the  $(N+1) \times (N+1)$  submatrix of  $S$  whose rows are  $\{1, \dots, N\} \cup \{i\}$  and columns are  $\{1, \dots, N\} \cup \{j\}$  with respect to the bases

$\mathcal{B}_1$  and  $\mathcal{B}_2$ . This submatrix has the following form:

$$\begin{bmatrix} 1 & 0 & \cdots & 0 & \lambda b_1 \\ 0 & 1 & \cdots & 0 & \lambda b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & \lambda b_N \\ \lambda a_1 & \lambda a_2 & \cdots & \lambda a_N & \lambda x_{ij} \end{bmatrix}_{(N+1) \times (N+1)}$$

for some scalars  $a_1, \dots, a_N$  and  $b_1, \dots, b_N$ . Since  $\text{rank}(S) \leq N$ , this determinant is 0. But by a direct computation, this implies that

$$\lambda x_{ij} = \lambda^2(a_1 b_1 + \cdots + a_N b_N)$$

and since  $\lambda$  is arbitrary, it follows that  $x_{ij} = 0$ . ■

The following proposition along with Proposition 3.1 gives the structure of closed subspaces of  $E \otimes E^*$ .

**PROPOSITION 3.2.** *Let  $E$  be a Banach space and  $M$  be a closed subspace of  $E \otimes E^*$ . Then there exists  $n_0 \in \mathbb{N}$  such that  $\text{rank}(T) \leq n_0$  for every  $T \in M$ .*

*Proof.* Let  $V_n = \{T \in E \otimes E^* : \text{rank}(T) \leq n\}$  for every  $n \in \mathbb{N}$ . We have  $M = \bigcup_{n \in \mathbb{N}} (M \cap V_n)$ . Since  $M$  is a Banach space, by the Baire category theorem there exists  $k_0$  such that  $\text{int}(M \cap V_{k_0}) \neq \emptyset$ . Let  $m \in M$  and  $\varepsilon > 0$  be such that  $B_M(m, \varepsilon) \subset V_{k_0}$ . Now  $B_M(m, \varepsilon) - B_M(m, \varepsilon) \subset V_{2k_0}$ . So  $B_M(0, \varepsilon) \subset B_M(m, \varepsilon) - B_M(m, \varepsilon) \subset V_{2k_0}$ , which implies that  $M \subset V_{2k_0}$ . Hence the ranks of the operators of  $M$  are uniformly bounded. ■

**REMARK 3.3.** We recall from [4] that  $\text{NA}(\mathcal{K}(\ell_2)) = \ell_2 \otimes \ell_2$ . Let  $Y$  be a closed subspace of  $\mathcal{K}(\ell_2)$  such that  $Y^\perp \subseteq \text{NA}(\mathcal{K}(\ell_2))$ . Now Propositions 3.1 and 3.2 imply that there exist  $f_1, \dots, f_N$  and  $e_1^*, \dots, e_N^*$  in  $\ell_2$  such that every  $T$  in  $Y^\perp$  can be written as

$$T = \sum_{i=1}^N e_i^* \otimes g_i + \sum_{j=1}^N b_j^* \otimes f_j$$

for some  $g_1, \dots, g_N$  and  $b_1^*, \dots, b_N^* \in \ell_2$ .

We now study proximality questions for factor reflexive subspaces of  $\mathcal{K}(\ell_2)$ . Let  $V$  be a finite-dimensional subspace of  $\ell_2$  and let

$$(1) \quad Z_V = \{S \in \mathcal{K}(\ell_2) : S(\ell_2) \subseteq V^\perp \text{ and } S^*(\ell_2) \subseteq V^\perp\}.$$

In other words, in an orthonormal basis  $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2$  where  $\mathcal{B}_1$  is a basis



of  $V$  and  $\mathcal{B}_2$  is a basis of  $V^\perp$ , the matrix of  $S$  has the form

$$\begin{bmatrix} [0]_{d \times d} & \vdots & 0 \\ \dots & \vdots & \dots \\ 0 & \vdots & [\alpha_{ij}] \end{bmatrix}$$

if and only if  $S \in Z_V$  where  $d$  is the dimension of  $V$ .

**PROPOSITION 3.4.** *For a finite-dimensional subspace  $V$  of  $\ell_2$  let  $Z_V$  be defined as in (1). Then  $Z_V$  is a proximal subspace of  $\mathcal{K}(\ell_2)$ .*

*Proof.* It suffices to show that every operator whose matrix relative to  $\mathcal{B}$  has the form

$$\begin{bmatrix} [\beta_{kl}^{(1)}]_{d \times d} & \vdots & [\beta_{mn}^{(2)}] \\ \dots & \vdots & \dots \\ [\beta_{pq}^{(3)}] & \vdots & 0 \end{bmatrix}$$

has a nearest point in  $Z_V$  (since we can translate by a vector in  $Z_V$ ). Such an operator has finite rank. Let

$$W = \text{span}\{V \cup S(V) \cup S^*(V)\}$$

and let  $W'$  be a finite-dimensional subspace of  $V^\perp$  such that

$$W \subseteq V \oplus W'.$$

Let  $\mathcal{B}' = \mathcal{B}_1 \cup \mathcal{B}'_2 \cup \mathcal{B}_3$  be an orthonormal basis of  $\ell_2$  such that  $\mathcal{B}_1$  (as before) is an orthonormal basis of  $V$ ,  $\mathcal{B}'_2$  is an orthonormal basis of  $W'$  and  $\mathcal{B}_3$  is an orthonormal basis of  $(V \oplus W')^\perp$ . The matrix of  $S$  relative to  $\mathcal{B}'$  is of the following form:

$$\begin{bmatrix} [\beta_{kl}^{(1)}]_{d \times d} & \vdots & [\beta_{mn}^{(2)}] & \vdots & 0 \\ \dots & \vdots & \dots & \vdots & \dots \\ [\beta_{pq}^{(3)}] & \vdots & [0]_{d' \times d'} & \vdots & 0 \\ \dots & \vdots & \dots & \vdots & \dots \\ 0 & \vdots & 0 & \vdots & 0 \\ \dots & \vdots & \dots & \vdots & \dots \end{bmatrix}$$

where  $d'$  is the dimension of  $W'$ . Let  $P : \ell_2 \rightarrow V \oplus W'$  be the orthogonal projection. If  $L \in Z_V$ , then  $P(S - L)P = S - PLP$  and we have  $PLP \in Z'$  with

$$Z' = \{L' \in \mathcal{K}(\ell_2) : L'(\ell_2) \subseteq W' \text{ and } L'^*(\ell_2) \subseteq W'\}.$$



Clearly  $Z'$  is a finite-dimensional vector subspace of  $Z_V$  consisting of operators whose matrix in  $\mathcal{B}'$  has the form

$$\begin{bmatrix} 0 & \vdots & 0 & \vdots & 0 \\ \dots & \vdots & \dots & \vdots & \dots \\ 0 & \vdots & [\gamma_{ij}] & \vdots & 0 \\ \dots & \vdots & \dots & \vdots & \dots \\ & \vdots & & \vdots & \\ 0 & \vdots & 0 & \vdots & 0 \\ & \vdots & & \vdots & \end{bmatrix}$$

Moreover since  $\|P\| = 1$ , we have

$$\|S - PLP\| = \|P(S - L)P\| \leq \|S - L\|.$$

Therefore

$$\inf\{\|S - L\| : L \in Z_V\} = \inf\{\|S - L'\| : L' \in Z'\},$$

and this infimum is attained since  $\dim(Z') < \infty$ , which completes the proof of the proposition. ■

We are now ready to state the main theorem of this section.

**THEOREM 3.5.** *Let  $Y$  be a closed subspace of  $\mathcal{K}(\ell_2)$  such that  $Y^\perp \subseteq \text{NA}(\mathcal{K}(\ell_2))$ . Then  $Y$  is a proximal subspace of  $\mathcal{K}(\ell_2)$ . In particular  $\mathcal{K}(\ell_2)$  is an  $\widetilde{R}(1)$  space.*

*Proof.* By Proposition 3.1, there is a finite-dimensional subspace  $V$  of  $\ell_2$  such that  $Z_V \subseteq Y$ . By Proposition 3.4, the space  $Z_V$  is a proximal subspace. Also  $\mathcal{K}(\ell_2)/Z_V$  is reflexive. Hence by Proposition 2.3,  $Y$  is a proximal subspace of  $\mathcal{K}(\ell_2)$ . ■

**REMARK 3.6.** If  $M \subseteq \text{NA}(\mathcal{K}(\ell_2)) \subseteq \mathcal{K}(\ell_2)^*$  is a norm-closed subspace, then  $M$  is necessarily reflexive. Indeed, since the dual unit ball of  $\mathcal{K}(\ell_2)$  is weakly sequentially complete,  $M^*$  is a quotient of  $X$  (see [1, Lemma 2.1]). Now being an  $M$ -embedded dual space,  $M^*$  and thus  $M$  is reflexive (see [5, Chapter III]).

We now prove that any  $\widetilde{R}(1)$  space with orthogonal linearity of norm attaining functionals is a  $\widetilde{P}$  space. For a proximal subspace  $Y$  of  $X$  let  $P_Y^{-1}(0) = \{x \in X : d(x, Y) = \|x\|\}$ .

**PROPOSITION 3.7.** *Let  $X$  be an  $\widetilde{R}(1)$  space such that  $\text{NA}(X)$  is orthogonally linear. Then  $X$  is a  $\widetilde{P}$  space.*

*Proof.* Let  $Z \stackrel{p}{\subseteq} Y \stackrel{p}{\subseteq} X$  be such that  $X/Z$  is reflexive. We have to show that  $Z \stackrel{p}{\subseteq} X$ . Since  $X$  is an  $\widetilde{R(1)}$  space, it suffices to show that  $Z^\perp \subseteq \text{NA}(X)$ .

The space  $Y^\perp$  is proximal in  $Z^\perp$  and thus  $Z^\perp = (P_{Y^\perp}^{-1}(0) \cap Z^\perp) + Y^\perp$ . We have  $Y \stackrel{p}{\subseteq} X$  and this implies that  $Y^\perp \subseteq \text{NA}(X)$ . Also  $Z \stackrel{p}{\subseteq} Y$  and so we have  $P_{Y^\perp}^{-1}(0) \cap Z^\perp \subseteq \text{NA}(X)$  and each functional in  $P_{Y^\perp}^{-1}(0) \cap Z^\perp$  is strongly orthogonal to  $Y^\perp$ . Since  $\text{NA}(X)$  is orthogonally linear this implies that  $Z^\perp \subseteq \text{NA}(X)$ . ■

REMARK 3.8. Since  $\text{NA}(\mathcal{K}(\ell_2))$  is a vector space, it follows by Proposition 3.7 that  $\mathcal{K}(\ell_2)$  is a  $\widetilde{P}$  space.

We next show that  $c_0$ -direct sums of reflexive spaces are  $\widetilde{R(1)}$  spaces.

LEMMA 3.9. Let  $\{X_i : i \in \mathbb{N}\}$  be a family of reflexive spaces and consider its  $c_0$ -direct sum  $X = (\bigoplus X_i)_{c_0}$ . Let  $M$  be a closed subspace of  $\text{NA}(X)$ . Then there exists a finite set  $A$  such that  $\text{supp}(f) \subseteq A$  for every  $f \in M$ .

*Proof.* Let  $V_n = \{f = (f_i) \in \text{NA}(X) : f_i = 0 \forall i > n_0\}$ . Then  $M = \bigcup_{n \in \mathbb{N}} (V_n \cap M)$ . Using the Baire category theorem arguments as in Proposition 3.2, we can get  $\varepsilon > 0$  and  $n_0$  such that  $B_M(0, \varepsilon) \subseteq V_{n_0}$ , which implies that  $M \subseteq V_{n_0}$  and this completes the proof. ■

It is easy to see that  $\text{NA}(X) = \{f = (f_i) \in X^* : f \text{ has only finitely many non-zero coordinates}\}$  and thus is a vector space.

PROPOSITION 3.10. Let  $\{X_i : i \in \mathbb{N}\}$  be a family of reflexive spaces and  $X = (\bigoplus X_i)_{c_0}$ . Let  $Y$  be a factor reflexive subspace of  $X$ . Then the following are equivalent.

- (i)  $Y$  is proximal in  $X$ .
- (ii)  $Y^\perp \subseteq \text{NA}(X)$ .
- (iii) there exists a finite set  $A$  such that  $\text{supp}(f) \subseteq A$  for every  $f \in Y^\perp$ .

*Proof.* (i)  $\Rightarrow$  (ii) by Lemma 2.2; (ii)  $\Rightarrow$  (iii) follows by 3.9; (iii)  $\Rightarrow$  (ii) is easy to see.

(ii)  $\Rightarrow$  (i). By Lemma 3.9 we can get  $n_0$  such that for all  $f = (f_i) \in Y^\perp$ ,  $f_i = 0$  if  $i > n_0$ . Let  $I = \{i : 1 \leq i \leq n_0\}$ ,

$$Y_1 = \left\{ x = (x_i) \in (\bigoplus_{\infty} X_i)_I : \sum_{i \in I} f_i(x_i) = 0 \forall f = (f_i) \in Y^\perp \right\}$$

and  $Y_2 = (\bigoplus_{c_0} X_i)_{N \setminus I}$ . Then clearly  $Y = Y_1 \oplus_{\infty} Y_2$  and  $Y_1$  is a closed subspace in a reflexive space  $(\bigoplus_{\infty} X_i)_I$ . So  $Y_1 \stackrel{p}{\subseteq} (\bigoplus_{\infty} X_i)_I$ . We have now  $Y = Y_1 \oplus_{\infty} Y_2 \stackrel{p}{\subseteq} X$ , which completes the proof. ■

THEOREM 3.11. Let  $\{X_i : i \in \mathbb{N}\}$  be a family of reflexive spaces and  $X = (\bigoplus X_i)_{c_0}$ . Then  $X$  is a  $\widetilde{P}$  space.

*Proof.* Proposition 3.10 implies that  $X$  is an  $\widetilde{R}(1)$  space and so by Proposition 3.7,  $X$  is a  $\widetilde{P}$  space (since  $\text{NA}(X)$  is a vector space). ■

REMARK 3.12. Let  $X$  be an  $\widetilde{R}(1)$  space such that  $\text{NA}(X)$  is a vector space. Let  $Y_1$  be a factor reflexive proximal subspace of  $X$  and  $Y_2$  be a finite-codimensional proximal subspace of  $X$ . Observe that  $Y_1^\perp$  is a reflexive subspace of  $\text{NA}(X)$  and  $Y_2^\perp$  is a finite-dimensional subspace of  $\text{NA}(X)$ . So  $Y_1^\perp + Y_2^\perp = (Y_1 \cap Y_2)^\perp \subseteq \text{NA}(X)$ . Since  $X$  is an  $\widetilde{R}(1)$  space, we conclude that  $Y_1 \cap Y_2$  is a factor reflexive proximal subspace of  $X$ .

REMARK 3.13. It is interesting to see whether the analogue of Lemma 2.13 holds true for factor reflexive spaces.

It follows from the discussion on  $\mathcal{K}(\ell_2)$  that if  $Y_1, \dots, Y_n$  are factor reflexive proximal subspaces of  $\mathcal{K}(\ell_2)$ , then  $Y_1 \cap \dots \cap Y_n$  is also proximal. Moreover, the following shows that for  $c_0$ -direct sums of reflexive spaces, the analogue of Lemma 2.13 holds true for factor reflexive spaces.

Let  $X$  be the  $c_0$ -direct sum of a family  $\{X_i : i \in \mathbb{N}\}$  of reflexive spaces. Let  $N$  be a closed subspace of  $\text{NA}(X)$ . Then there is a finite set  $A$  of  $\mathbb{N}$  such that  $N \subseteq M = (\bigoplus_{\ell^1} X_i^*)_{i \in A}$ . But  $M$  is a reflexive space. Hence so is  $N$ . Now by Propositions 3.10 and 2.3,  $N_\perp$  is proximal in  $X$ .

Let  $Y_1$  and  $Y_2$  be two factor reflexive proximal subspaces of  $X$ . As before there exist finite subsets  $A_1$  and  $A_2$  of  $\mathbb{N}$  such that  $Y_1^\perp \subseteq M_1 = (\bigoplus_{\ell^1} X_i^*)_{i \in A_1}$  and  $Y_2^\perp \subseteq M_2 = (\bigoplus_{\ell^1} X_i^*)_{i \in A_2}$ . Now by duality  $(M_1 \cap M_2)_\perp \subseteq Y_1 \cap Y_2 \subseteq (\bigoplus_{c_0} X_i)_{i \in \mathbb{N} \setminus (A_1 \cup A_2)}$ . But  $(M_1 \cap M_2)_\perp$  is proximal in  $X$ . Thus by Proposition 2.3 again,  $Y_1 \cap Y_2$  is proximal in  $X$ .

We conclude this section with the following questions.

- (i) Is  $X$  a  $\widetilde{P}$  space only if it is an  $\widetilde{R}(1)$  space and  $\text{NA}(X)$  is orthogonally linear?
- (ii) Is there any example of an  $R(1)$  space  $X$  and  $Y \subset X$  such that the quotient is infinite-dimensional and reflexive, every finite-codimensional subspace containing  $Y$  is proximal in  $X$ , but  $Y$  itself is not proximal in  $X$ ?
- (iii) We do not know whether  $\mathcal{K}(\ell_p)$  for  $1 < p < \infty$  and  $p \neq 2$  is at least a  $P$  space.

4. Renorming of  $\widetilde{R}(1)$  spaces. It is known that given a separable space there is an equivalent smooth norm with the same set of norm attaining functionals, i.e., proximal hyperplanes are the same (see [2]). A natural question then is to know whether proximal factor reflexive subspaces remain the same. In this section, we answer this question affirmatively. We start with a crucial and simple lemma which applies in particular to all separable spaces.

**LEMMA 4.1.** *Let  $(X, \|\cdot\|)$  be a normed linear space. Let  $L$  be any weakly compact convex symmetric subset of  $X$ . Let  $\|\cdot\|$  be the norm whose unit ball satisfies  $B_X(\|\cdot\|) = B_X(\|\cdot\|) + L$ . Let  $Y$  be a closed subspace of  $(X, \|\cdot\|)$ . If  $Y$  is proximal in  $(X, \|\cdot\|)$  then  $Y$  is proximal in  $(X, \|\cdot\|)$ .*

*Proof.* Let  $x \in X$  be such that  $d_{\|\cdot\|}(x, Y) = 1$ . Then for every  $n \in \mathbb{N}$ , we have  $Y \cap (B_{(X, \|\cdot\|)}(x, 1 + 1/n) + (1 + 1/n)L) \neq \emptyset$ . Let  $y_n = t_n + l_n \in Y \cap (B_{(X, \|\cdot\|)}(x, 1 + 1/n) + (1 + 1/n)L)$ , where  $y_n \in Y$  and  $l_n \in L$ . Let  $\{l_{n_i}\}$  be a weakly converging subsequence of  $\{l_n\}$  and let  $x + l = w\text{-}\lim(x + l_{n_i})$ . We have  $d_{\|\cdot\|}(x + l, Y) = 1$ . Since  $Y$  is proximal in  $(X, \|\cdot\|)$ , we have  $d_{\|\cdot\|}(x + l, Y) = \|x + l - y\| = 1$  for some  $y \in Y$ . If  $v = x + l - y$ , one has  $x + l - v = y \in Y$  and thus  $d_{\|\cdot\|}(x, Y) = 1 = \|x - y\|$  and  $Y$  is proximal in  $(X, \|\cdot\|)$ . ■

We now prove the main theorem of this section which shows that a separable  $\widetilde{R}(1)$  space can be smoothly renormed preserving its proximality properties. In particular these arguments also hold for  $R(1)$  spaces.

**THEOREM 4.2.** *Let  $(X, \|\cdot\|)$  be a separable  $\widetilde{R}(1)$  space. Then there exists an equivalent Gateaux smooth norm  $\|\cdot\|$  on  $X$  such that  $X$  with this new norm is again  $\widetilde{R}(1)$ .*

*Proof.* By Theorem 9(iv) from [2] there exists an equivalent Gateaux smooth norm  $\|\cdot\|$  on  $X$  such that  $\text{NA}((X, \|\cdot\|)) = \text{NA}((X, \|\cdot\|))$ . Indeed, let  $\{x_n\}$  be a dense subset of  $B_X$ , define  $T: \ell_2 \rightarrow X$  by  $T(\alpha) = \sum_{n=1}^{\infty} 2^{-n} \alpha_n x_n$ , and let  $K = T(B_{\ell_2})$ . The set  $K$  is convex, symmetric and norm compact. Let  $\|\cdot\|$  be the norm whose unit ball satisfies  $B_X(\|\cdot\|) = B_X(\|\cdot\|) \div K$ . Let  $X = (X, \|\cdot\|)$  and  $X_1 = (X, \|\cdot\|)$ . By Lemma 4.1,  $f \in \text{NA}(X)$  if and only if  $f \in \text{NA}(X_1)$ . Moreover, for  $f \in X^*$ ,

$$\begin{aligned} (2) \quad \|f\|^* &= \sup\{|f(x_1)| : x_1 \in B_{X_1}\} \\ &= \sup\{|f(x+k)| : x \in B_X, k \in K\} \\ &= \sup\{|f(x)| : x \in B_X\} + \sup\{|f(k)| : k \in K\} \\ &= \|f\|^* + \sup\{|f(T(\alpha))| : \alpha \in B_{\ell_2}\} = \|f\|^* + \|T^*(f)\|_2. \end{aligned}$$

Since  $T^*$  is one-to-one and  $\|\cdot\|_2$  is strictly convex, it follows that  $\|\cdot\|^*$  is strictly convex and thus  $\|\cdot\|$  is Gateaux smooth.

Let  $Y$  be a factor reflexive subspace of  $X$ . Suppose that  $Y^\perp \subseteq \text{NA}(X) = \text{NA}(X_1)$ . Since  $X$  is an  $\widetilde{R}(1)$  space,  $Y$  is proximal in  $X$ . Let  $Y_1 = (Y, \|\cdot\|)$ . Then by Lemma 4.1,  $Y$  is proximal in  $(X, \|\cdot\|)$ , which completes the proof. ■

**REMARK 4.3.** By the above results,  $c_0$  and more generally  $c_0$ -direct sums of sequences of reflexive spaces admit Gateaux smooth norms such that with these new norms these spaces are still  $\widetilde{R}(1)$  spaces.

**5. Linearity of  $\text{NA}(Y)$  for a hyperplane  $Y$  in  $c_0$ .** We first recall that  $\text{NA}(c_0)$  is a vector space and for any proximal hyperplane  $Y$  in  $c_0$ ,  $\text{NA}(Y)$  is a vector space (by Proposition 2.5). However when  $Y$  is not proximal,  $\text{NA}(Y)$  can fail to be linear. In this direction we present an example which shows that if  $f = (f_i) \in \ell_1$  is not norm attaining then  $\text{NA}(\ker f)$  need not even be orthogonally linear.

**EXAMPLE 5.1.** Let  $f = (1/2, 1/2, 1/4, 1/8, \dots) \in \ell_1$ . Let  $X = \ker f$ . It can be easily seen that  $\text{NA}(X)$  is not a vector space ([3]). We show that it is not even orthogonally linear. Indeed, let  $g = (1, 0, 0, \dots)$  and  $H = (0, 0, 1, 0, 0, \dots)$ . Now  $x = (0, -1/2, 1, 0, 0, \dots) \in S_{\ker g}$  is such that  $H(x) = \|H\| = \|H|_{\ker g \cap X}\| = 1$ . So  $H$  is strongly orthogonal to  $g$  in  $X^*$ . But  $g + H = (1, 0, 1, 0, 0, \dots)$  and  $\|g + H\|_{X^*} = 2 = 1 + \sum_{i=1}^{\infty} 2^{-i}$ . Let  $x^{(n)} = (1, -1, \sum_{i=1}^n 2^{-i}, -1, \dots, -1_{n+4}, 0, \dots)$ , where  $1_i = 1$  for  $4 \leq i \leq n + 4$ . Then  $x^{(n)}$  is in  $B_X$  and  $(g + H)(x^{(n)}) \rightarrow 2$  but there is no  $x \in B_X$  such that  $(g + H)(x) = 2$ ; this implies that  $g + H \notin \text{NA}(X)$ . Hence  $\text{NA}(X)$  is not orthogonally linear. Thus by Theorem 3 of [3] and Corollary 5 of [7],  $X$  is an  $R(1)$ -space but not a  $P$  space.

In view of the above example, one can ask the following questions.

**QUESTION 5.2.** *Are there any non-proximal hyperplanes of  $c_0$  such that the set of all norm attaining functionals is a vector space?*

**QUESTION 5.3.** *Do linearity and orthogonal linearity coincide in hyperplanes of  $c_0$ ? This is a particular case of Question 1 from [7].*

We answer affirmatively the above questions.

To state the next result we need the following notation.

Let  $f = (f_i) \in S_{\ell_1}$ . Suppose  $f \notin \text{NA}(c_0)$ . Let  $|f_{i_1}| = \sup\{|f_i| : i \in \mathbb{N}\}$  and  $|f_{i_j}| = \sup\{|f_i| : i \in \mathbb{N} \setminus \{i_1, \dots, i_{j-1}\}\}$  for  $j \geq 2$ . Then  $\{|f_{i_n}|\}$  is a decreasing sequence. Let  $Y = \ker f$ .

**PROPOSITION 5.4.** *Suppose  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$ . Then  $Y$  is isometric to  $c_0$  and thus  $\text{NA}(Y)$  is a vector space. Moreover  $\text{NA}(Y) = \{g|_Y : g \in \text{NA}(c_0) \text{ with the } i_1 \text{th coordinate zero}\}$ .*

*Proof.* Let  $y = (y_i) \in Y$  and let  $T : Y \rightarrow c_0(\mathbb{N} \setminus \{i_1\})$  be defined by  $T(y) = (y_i)_{i \in \mathbb{N} \setminus \{i_1\}}$ . We have  $\|T(y)\|_{\infty} = \|y\|_{\infty}$  and  $T$  is onto  $c_0(\mathbb{N} \setminus \{i_1\})$ . Thus we have

$$\begin{aligned} \text{NA}(Y) &= T^*(\text{NA}(c_0(\mathbb{N} \setminus \{i_1\}))) \\ &= \{g|_Y : g \in \text{NA}(c_0(\mathbb{N})) \text{ with the } i_1 \text{th coordinate zero}\}. \blacksquare \end{aligned}$$

First we prove the converse for a particular hyperplane. Let  $f = (f_i) \in S_{\ell_1} \setminus \text{NA}(c_0)$  be such that each  $f_i$  has a constant sign for  $i \in \mathbb{N}$ . As above,

let  $|f_{i_1}| = \max\{|f_i| : i \in \mathbb{N}\}$  and  $|f_{i_j}| = \sup\{|f_i| : i \in \mathbb{N} \setminus \{i_1, i_2, \dots, i_{j-1}\}\}$  for  $j \geq 2$ . Let  $Y = \ker f$ .

**PROPOSITION 5.5.** *If  $NA(Y)$  is a vector space then  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$ .*

*Proof.* Suppose  $NA(Y)$  is a vector space. We argue by contradiction. Assume that there exists a finite subset  $J_1$  of  $\mathbb{N} \setminus \{i_1\}$  such that  $|f_{i_1}| \leq \sum_{i \in J_1} |f_i|$ . Then there exist  $\alpha^{(1)} = (\alpha_i^{(1)})$  in  $[-1, 1]^{|J_1|}$  and  $\alpha^{(2)} = (\alpha_j^{(2)})$  in  $[-1, 1]^{|(J_1 \cup \{i_1\}) \setminus \{i_2\}|}$  such that

$$-f_{i_1} = \sum_{i \in J_1} \alpha_i^{(1)} f_i \quad \text{and} \quad -f_{i_2} = \sum_{j \in (J_1 \cup \{i_1\}) \setminus \{i_2\}} \alpha_j^{(2)} f_j.$$

Let  $g_1 = e_{i_1}$  and  $g_2 = e_{i_2}$ . It is easy to see that  $g_1|_Y, g_2|_Y \in NA(Y)$ . Indeed, let  $y^{(1)} = (y_i^{(1)})$  and  $y^{(2)} = (y_i^{(2)})$  in  $S_Y$ , where

$$y_i^{(1)} = \begin{cases} \alpha_i^{(1)} & \text{if } i \in J_1, \\ 1 & \text{if } i = i_1, \\ 0 & \text{otherwise,} \end{cases} \quad y_i^{(2)} = \begin{cases} \alpha_i^{(2)} & \text{if } i \in (J_1 \cup \{i_1\}) \setminus \{i_2\}, \\ 1 & \text{if } i = i_2, \\ 0 & \text{otherwise.} \end{cases}$$

Then  $g_1|_Y(y^{(1)}) = 1 = \|g_1|_Y\|_{Y^*}$  and  $g_2|_Y(y^{(2)}) = 1 = \|g_2|_Y\|_{Y^*}$ . We now have

**LEMMA 5.6.** *The following are equivalent.*

- (i)  $g_1|_Y + g_2|_Y \in NA(Y)$ .
- (ii) *There exists a finite subset  $J_2$  of  $\mathbb{N} \setminus \{i_1, i_2\}$  such that  $|f_{i_1}| + |f_{i_2}| \leq \sum_{i \in J_2} |f_i|$ .*

*Proof of Lemma 5.6.* (i)  $\Rightarrow$  (ii). Suppose  $g_1|_Y + g_2|_Y \in NA(Y)$  but there is no finite subset  $J_2$  of  $\mathbb{N} \setminus \{i_1, i_2\}$  such that  $|f_{i_1}| + |f_{i_2}| \leq \sum_{i \in J_2} |f_i|$ . Let  $y = (y_i) \in S_Y$  be such that  $(g_1 + g_2)(y) = |y_{i_1} + y_{i_2}| = \|(g_1 + g_2)|_Y\|_{Y^*}$ . It is easy to see that  $y_{i_1}$  and  $y_{i_2}$  have the same sign. We have  $f(y) = 0$ , so  $-(y_{i_1} f_{i_1} + y_{i_2} f_{i_2}) = \sum_{i=1, i \neq i_1, i_2}^{\infty} y_i f_i$ , which implies that

$$|y_{i_1} f_{i_1} + y_{i_2} f_{i_2}| = \left| \sum_{i \neq i_1, i_2} y_i f_i \right| \leq \sum_{i \neq i_1, i_2} |y_i f_i| < \sum_{i \neq i_1, i_2} |f_i|.$$

Let  $\alpha_{i_1}, \alpha_{i_2} \in [-1, 1]$  be such that  $\text{sign}(\alpha_{i_1}) = \text{sign}(\alpha_{i_2}) = \text{sign}(y_{i_1}) = \text{sign}(y_{i_2})$ ,  $|y_{i_1}| < |\alpha_{i_1}|$ ,  $|y_{i_2}| < |\alpha_{i_2}|$  and

$$-(\alpha_{i_1} f_{i_1} + \alpha_{i_2} f_{i_2}) = \sum_{i=1, i \neq i_1, i_2}^{\infty} |f_i|.$$

Let  $\alpha_{i_1}^{(n)}$  and  $\alpha_{i_2}^{(n)}$  in  $[-1, 1]$  be such that

$$-(\alpha_{i_1}^{(n)} f_{i_1} + \alpha_{i_2}^{(n)} f_{i_2}) = \sum_{i=1, i \neq i_1, i_2}^n |f_i|,$$

$\alpha_{i_1}^{(n)} \rightarrow \alpha_{i_1}$  and  $\alpha_{i_2}^{(n)} \rightarrow \alpha_{i_2}$ . Now let  $y^{(n)} = (y_i^{(n)})$ , where

$$y_i^{(n)} = \begin{cases} -\text{sign}(f_i) & \text{if } i \in \{1, \dots, n\} \setminus \{i_1, i_2\}, \\ \alpha_{i_1}^{(n)} & \text{if } i = i_1, \\ \alpha_{i_2}^{(n)} & \text{if } i = i_2. \end{cases}$$

Then  $(g_1 + g_2)(y^{(n)}) = \alpha_{i_1}^{(n)} + \alpha_{i_2}^{(n)}$  and  $(g_1 + g_2)(y^{(n)}) \rightarrow \alpha_{i_1} + \alpha_{i_2}$ . This contradicts the fact that  $\|(g_1 + g_2)|_Y\| = |y_{i_1} + y_{i_2}|$ . So there exists a finite subset  $J_2$  of  $\mathbb{N} \setminus \{i_1, i_2\}$  such that  $|f_{i_1}| + |f_{i_2}| \leq \sum_{i \in J_2} |f_i|$ .

(ii)  $\Rightarrow$  (i). Assume there exists a finite subset  $J_2$  of  $\mathbb{N} \setminus \{i_1, i_2\}$  such that  $|f_{i_1}| + |f_{i_2}| \leq \sum_{i \in J_2} |f_i|$ . Then there exists  $\alpha_i \in [-1, 1]^{|J_2|}$  such that  $|f_{i_1}| + |f_{i_2}| = -\sum_{i \in J_2} \alpha_i f_i$ . Consider  $y = (y_i)$ , where

$$y_i = \begin{cases} \alpha_i & \text{if } i \in J_2, \\ \text{sign}(f_{i_1}) & \text{if } i = i_1, \\ \text{sign}(f_{i_2}) & \text{if } i = i_2, \\ 0 & \text{otherwise.} \end{cases}$$

Then  $|(g_1 + g_2)(y)| = 2$  and so  $g_1|_Y + g_2|_Y \in \text{NA}(Y)$ .

*End of proof of Proposition 5.5.* If  $g_1 + g_2 \notin \text{NA}(Y)$  we are done. Otherwise consider  $g_3 = e_{i_3}$ . Then as in Lemma 5.6 we can show that  $g_1 + g_2 + g_3 \in \text{NA}(Y)$  if and only if there exists a finite subset  $J_3$  of  $\mathbb{N} \setminus \{i_1, i_2, i_3\}$  such that  $|f_{i_1}| + |f_{i_2}| + |f_{i_3}| \leq \sum_{i \in J_3} |f_i|$ . Since  $f \in S_{\ell_1}$ , there exists  $n_0$  such that  $\sum_{j=1}^{n_0} |f_{i_j}| \geq 2/3$ . So this process has to stop, and we get  $n < n_0$  such that  $\sum_{j=1}^n g_j$  and  $g_{n+1}$  are in  $\text{NA}(Y)$  but  $\sum_{j=1}^{n+1} g_j$  is not, contrary to the assumption that  $\text{NA}(Y)$  is a vector space.

**REMARK 5.7.** Lemma 5.6 is not true if  $f_i$ 's do not have constant sign. Indeed, let  $f = (1, -1, 1/2, 1/4, 1/8, \dots)$ . Then both  $e_1 = (1, 0, 0, \dots)$  and  $e_2 = (0, 1, 0, 0, \dots)$  are in  $\text{NA}(\ker f)$  and also  $e_1 + e_2 \in \text{NA}(\ker f)$  but Lemma 5.6(ii) is not satisfied. But here  $e_1 + e_2 + e_3 \notin \text{NA}(\ker f)$ .

As usual let  $f = (f_i) \in S_{\ell_1} \setminus \text{NA}(c_0)$ . Let  $|f_{i_1}| = \max\{|f_i| : i \in \mathbb{N}\}$  and  $|f_{i_j}| = \max\{|f_i| : i \in \mathbb{N} \setminus \{i_1, \dots, i_{j-1}\}\}$  for  $j \geq 2$ . Let  $Y = \ker f$ . Then we have

**THEOREM 5.8.**  $\text{NA}(Y)$  is a vector space if and only if  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$ . Moreover if  $\text{NA}(Y)$  is a vector space, then  $\text{NA}(Y) = \{h|_Y : h \in \text{NA}(c_0) \text{ with the } i_1 \text{th coordinate zero}\}$ .

*Proof.* Let  $f = (f_i) \in S_{\ell_1}$ ,  $|f| = (|f_i|)$  and let

$$\text{sign}(f_i) = \begin{cases} 1 & \text{if } f_i \geq 0, \\ -1 & \text{if } f_i < 0. \end{cases}$$

Now we define a map  $T : c_0 \rightarrow c_0$  by  $T(x) = (\text{sign}(f_i)x_i)$ . Then  $T$  is an invertible isometry and  $T(\ker f) = \ker |f|$ . Hence  $\text{NA}(\ker |f|) = T^*(\text{NA}(\ker f))$ .



If  $\text{NA}(\ker f)$  is a vector space, then so is  $\text{NA}(\ker |f|)$ . By Proposition 5.5,  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$ . The converse follows again by Proposition 5.4. The second part is a consequence of Proposition 5.4. ■

**THEOREM 5.9.** *Let  $f = (f_i) \in S_{\ell_1}$ . Then  $\text{NA}(\ker f)$  is orthogonally linear if and only if it is linear.*

*Proof.* Suppose  $\text{NA}(\ker f)$  is orthogonally linear. Let  $T$  be an isometry from  $c_0$  to  $c_0$  defined by  $T(x) = (\text{sign}(f_i)x_i)$  as in the previous proof. Then  $\text{NA}(\ker f)$  is orthogonally linear if and only if  $\text{NA}(\ker |f|)$  is. Now it is enough to prove that, if  $\text{NA}(\ker |f|)$  is orthogonally linear, then  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$  where  $|f_{i_j}| = \sup\{|f_i| : i \in \mathbb{N} \setminus \{i_1, \dots, i_{j-1}\}\}$  for  $j \geq 1$  and  $i_0 = \{0\}$ . Suppose not. Let  $g_j = e_{i_j}$  for  $j \geq 1$ . Then  $g_j \in \text{NA}(\ker |f|)$  for  $j \geq 1$ . It is easy to see that  $g_3$  is strongly orthogonal to  $g_1$ . Thus  $g_1 + g_3 \in \text{NA}(\ker |f|)$  by orthogonal linearity. Now as in the proof of Lemma 5.6, there exists a finite subset  $J_2 \subset \mathbb{N} \setminus \{i_1, i_3\}$  such that  $|f_{i_1}| + |f_{i_3}| \leq \sum_{i \in J_2} |f_i|$ . There exists  $\alpha_i \in [-1, 1]$  for  $i \in J_2 \cup \{i_2, i_4\}$  such that

$$-(|f_{i_1}| - |f_{i_3}|) = \sum_{i \in J_2 \cup \{i_2, i_4\}} \alpha_i |f_i| \quad \text{and} \quad \alpha_{i_4} \in \{-1, 1\}.$$

Now let  $y = (y_i)$ , where

$$y_i = \begin{cases} \alpha_i & \text{if } i \in J_2 \cup \{i_2, i_4\}, \\ 1 & \text{if } i = i_1, \\ -1 & \text{if } i = i_3, \\ 0 & \text{otherwise.} \end{cases}$$

Then  $y \in S_{\ker(g_1+g_3)}$  and  $|g_4(y)| = 1$ , which implies that  $g_4$  is strongly orthogonal to  $g_1 + g_3$ . Thus  $g_1 + g_3 + g_4 \in \text{NA}(X)$  by orthogonal linearity. Proceeding as in Proposition 5.5, we show that there exists  $n \in \mathbb{N}$  such that  $g_{n+1}$  is strongly orthogonal to  $g_1 + g_3 + g_4 + \dots + g_n$  but  $g_1 + g_3 + g_4 + \dots + g_{n+1}$  is not in  $\text{NA}(X)$ , which contradicts the orthogonal linearity of  $\text{NA}(X)$ . The converse is trivial. ■

**COROLLARY 5.10.** *Let  $Y$  be a non-proximinal hyperplane in  $c_0$ . Let  $f = (f_i) \in \ell_1$  be such that  $Y = \ker f$ . Then the following are equivalent.*

- (i)  $Y$  is a  $P$  space.
- (ii)  $|f_{i_1}| \geq \sum_{i=1, i \neq i_1}^{\infty} |f_i|$ , where  $|f_{i_1}| = \max\{|f_i| : i \in \mathbb{N}\}$  and  $|f_{i_j}| = \sup\{|f_i| : i \in \mathbb{N} \setminus \{i_1, \dots, i_{j-1}\}\}$  for  $j \geq 2$ .
- (iii)  $\text{NA}(Y) = \{h|_Y : h \in \text{NA}(c_0) \text{ with the } i_1 \text{th coordinate zero}\}$ .

*Proof.* (ii)  $\Leftrightarrow$  (iii) follows from Theorem 5.8; (i)  $\Rightarrow$  (iii) follows by Corollary 5 of [7], Theorem 5.9 and Theorem 5.8; (iii)  $\Rightarrow$  (i) follows by Theorem 3 of [3] and Corollary 5 of [7]. ■

## REFERENCES

- [1] P. Bandyopadhyay and G. Godefroy, *Linear structures in the set of norm-attaining functionals of a Banach space*, preprint, 2005.
- [2] G. Debs, G. Godefroy and J. Saint-Raymond, *Topological properties of the set of norm-attaining linear functionals*, *Canad. J. Math.* 47 (1995), 318–329.
- [3] G. Godefroy and V. Indumathi, *Proximality in subspaces of  $c_0$* , *J. Approx. Theory* 101 (1999), 175–181.
- [4] G. Godefroy, V. Indumathi and F. Lust Piquard, *Strong subdifferentiability of convex functionals and proximality*, *ibid.* 116 (2002), 397–415.
- [5] P. Harmand, D. Werner and W. Werner, *M-Ideals in Banach Spaces and Banach Algebras*, *Lecture Notes in Math.* 1547, Springer, Berlin, 1993.
- [6] V. Indumathi, *Proximinal subspaces of finite codimension in general normed linear spaces*, *Proc. London Math. Soc.* 45 (1982), 435–455.
- [7] —, *On transitivity of proximality*, *J. Approx. Theory* 49 (1987), 130–143.
- [8] D. Narayana and T. S. S. R. K. Rao, *Proximality in generalized direct sums*, *Int. J. Math. Math. Sci.* 67 (2004), 3663–3670.
- [9] W. Pollul, *Reflexivität und Existenz-Teilräume in der linearen Approximationstheorie*, *Gesellschaft für Mathematik und Datenverarbeitung, Bonn, Ber.* 53, 1972.
- [10] I. Singer, *Best Approximations in Normed Linear Spaces by Elements of Linear Subspaces*, *Grundlehren Math. Wiss.* 171, Springer, Berlin, 1970.

Department of Mathematics  
Indian Institute of Science  
Bangalore 560012, India  
E-mail: narayana@math.iisc.ernet.in

Stat-Math Unit  
Indian Statistical Institute  
R. V. College P.O.  
Bangalore 560059, India  
E-mail: tas@isibang.ac.in

Received 9 February 2004;  
revised 20 April 2005

(4421)