

A note on equivariant Euler characteristic

AMIYA MUKHERJEE and ANIRUDDHA C NAOLEKAR

Stat-Math Division, Indian Statistical Institute, Calcutta 700 035, India

E-mail: amiya@isical.ernet.in

MS received 2 July 1996

Abstract. We give a new equivariant cohomological characterization of the equivariant Euler characteristic of a G -simplicial set as defined by Brown. This implies in particular that the equivariant Euler characteristic is a G -homotopy invariant.

Keywords. G -simplicial set; equivariant cohomology; Euler characteristic.

1. Introduction

Serre [5] and Brown [2] defined the Euler characteristic of a discrete group G of finite homological type. Recall that the cohomological dimension $\text{cd } G$ is $\inf n$ so that the $\mathbb{Z}G$ -module \mathbb{Z} with trivial G -action admits projective resolutions of length $\leq n$, and that G has finite virtual cohomological dimension $\text{vcd } G$ if there exists a finite index subgroup with finite cohomological dimension. A group G is of finite homological type if

- (1) $\text{vcd } G < \infty$, and
- (2) every torsion-free subgroup of finite index has finitely generated rational homology.

If the rational homology of G is finitely generated, then the 'naive' Euler characteristic of G is the integer

$$\tilde{\chi}(G) = \sum (-1)^i \dim_{\mathbb{Q}} H_i(G; \mathbb{Q}),$$

and if G is of finite homological type, then its Euler characteristic is the rational number

$$\chi(G) = \tilde{\chi}(H)/[G:H],$$

where H is a torsion-free subgroup of finite index $[G:H]$. It is a result of Brown [2] that $\chi(G)$ is independent of the choice of H .

Next, if K is a G -simplicial set, where G is discrete, such that the simplicial set K/G has finitely many non-degenerate cells, and each isotropy subgroup G_x of a simplex x of K is of finite homological type, then the equivariant Euler characteristic of K is

$$\chi_G(K) = \sum (-1)^{\dim x} \chi(G_x),$$

where the sum is over the representatives of non-degenerate simplexes of K/G .

Now suppose that O_G denotes the category of canonical orbits of G whose objects are left coset spaces G/H and whose morphisms are G -maps $\hat{g}: G/H \rightarrow G/H'$ coming from a subconjugacy relation $g^{-1}Hg \leq H'$. Let $\lambda_{\mathbb{Q}}$ denote the contravariant functor from O_G to the category of vector spaces over \mathbb{Q} such that $\lambda_{\mathbb{Q}}(G/H) = \text{Hom}(\mathbb{Q}(G/H), \mathbb{Q})$ where $\mathbb{Q}(G/H)$ is the vector space over \mathbb{Q} with basis G/H , and $\lambda_{\mathbb{Q}}(\hat{g}) = \text{Hom}(\mathbb{Q}(\hat{g}), id)$.

In § 2, we define equivariant cohomology $H_G^*(K; \lambda_{\mathbb{Q}})$ of K which is G -homotopy invariant. In fact, this cohomology is the simplicial analogue of the Bredon cohomology [1]. The purpose of the present paper is to prove the following theorems.

Theorem 1. *If the action of a discrete group G on a G -simplicial set K is such that (i) K/G has only finitely many non-degenerate cells, and (ii) every isotropy subgroup of every simplex of K has finite index in G , and if $\lambda_{\mathbb{Q}}$ is as defined above, then the cohomology groups $H_G^*(K; \lambda_{\mathbb{Q}})$ are finitely generated.*

Theorem 2. *If G acts freely on a simplicial set K satisfying the conditions (i) and (ii) of Theorem 1 then $\chi_G(K) = \chi(G) \sum (-1)^i \dim_{\mathbb{Q}} H_G^i(K; \lambda_{\mathbb{Q}})$, where the summation is from $i = 0$ to $i = \dim K/G$.*

Theorem 3. *If G is of finite cohomological dimension and finite homological type, and K is a G -simplicial set satisfying the conditions of the above theorem, then*

$$\chi_G(K) = \chi(G) \sum (-1)^i \dim_{\mathbb{Q}} H_G^i(K; \lambda_{\mathbb{Q}}),$$

where the summation is from $i = 0$ to $i = \dim K/G$.

Thus $\chi_G(K)$ is a G -homotopy invariant. In particular, if G is free of rank n , then it is of finite homological type because its virtual cohomological dimension is n . Moreover its rational homology is finitely generated because there exists a $K(G, 1)$ with one 0-cell and n 1-cells. Therefore $\chi(G) = 1 - n$, and

$$\chi_G(K) = (1 - n) \sum (-1)^i \dim_{\mathbb{Q}} H_G^i(K; \lambda_{\mathbb{Q}}),$$

if K is as in Theorem 1.

The plan of the paper is as follows. In § 2, we discuss some basic results with only sketches of proofs, the details of which may be worked out without difficulty. The proofs of the theorems appear in § 3.

2. Equivariant cohomology of a G -simplicial set

A G -simplicial set is a simplicial set K together with an action of G by simplicial maps which commute with the face and degeneracy maps d_i and s_i .

We define the equivariant cohomology $H_G^*(K; \lambda)$ of K with a coefficient system λ (which is a contravariant functor from O_G to the category $R\text{-mod}$ of R -modules, R being a commutative ring with (1) as follows. Let $C^n(K; \lambda)$ be the R -module of functions c defined on n -simplexes x of K such that $c(x) \in \lambda(G/G_x)$, where G_x is the isotropy subgroup of x in G . Then define coboundary $\delta: C^n(K; \lambda) \rightarrow C^{n+1}(K; \lambda)$ by

$$\delta(c)(x) = \sum_{i=0}^{n+1} (-1)^i \lambda(d_i x \rightarrow x) c(d_i x),$$

where $\lambda(d_i x \rightarrow x)$ is the homomorphism $\lambda(G/G_{d_i x}) \rightarrow \lambda(G/G_x)$ induced from the G -map $G/G_x \rightarrow G/G_{d_i x}$ given by the inclusion $G_x \subseteq G_{d_i x}$.

We define an action of G on $C^n(K; \lambda)$ by $(gc)(x) = \lambda(\hat{g})(c(g^{-1}x))$ where $\lambda(\hat{g})$ is the isomorphism $\lambda(G/G_{g^{-1}x}) \rightarrow \lambda(G/G_x)$ induced by the conjugacy relation

$g^{-1}G_xg = G_{g^{-1}x}$. Let $C_G^n(K; \lambda)$ be the submodule of G -invariant cochains $(C^n(K; \lambda))^G$. Clearly this makes $C_G^*(K; \lambda)$ a cochain complex, and so we may define

$$H_G^n(K; \lambda) = H_n(C_G^*(K; \lambda)).$$

Note that if the action of G on K is free, then we have

$$H_G^*(K; \lambda) \cong H^*(K/G; \lambda(G/\{e\}))$$

for every coefficient system λ .

Clearly a G -simplicial map $f: K \rightarrow L$ induces a cochain map $f^\#: C_G^*(L; \lambda) \rightarrow C_G^*(K; \lambda)$ defined by $f^\#(c)(x) = \lambda(fx \rightarrow x)c(fx)$, where $\lambda(fx \rightarrow x): \lambda(G/G_{fx}) \rightarrow \lambda(G/G_x)$ is the homomorphism induced by the inclusion $G_x \subseteq G_{fx}$. Then $f^\#$ induces homomorphism $f^*: H_G^*(L; \lambda) \rightarrow H_G^*(K; \lambda)$ satisfying the usual functorial properties.

Lemma 4. If $f, g: K \rightarrow L$ are G -homotopic G -simplicial maps, then

$$f^* = g^*: H_G^*(L; \lambda) \rightarrow H_G^*(K; \lambda).$$

Sketch of Proof. The cochain maps $f^\#, g^\#: C_G^*(L; \lambda) \rightarrow C_G^*(K; \lambda)$ are cochain homotopic by $h: C_G^n(L; \lambda) \rightarrow C_G^{n-1}(K; \lambda)$ given by

$$h(c)(x) = \sum_{j=0}^{n-1} (-1)^j \lambda(h_jx \rightarrow x)c(h_jx),$$

where $h_j: K_n \rightarrow L_{n+1}$ are G -functions constituting a G -homotopy from f to g . ■

Alternatively, the cochain complex $C_G^*(K; \lambda)$ may be defined as follows. Consider for each $n = \geq 0$ a coefficient system $\underline{C}_n(K): O_G \rightarrow R\text{-mod}$ by setting $\underline{C}_n(K)(G/H) = C_n(K^H; R)$ which is the free R -module generated by the n -simplexes of K^H , and, for a G -map $\hat{g}: G/H \rightarrow G/H', g^{-1}Hg \subseteq H'$, setting $\underline{C}_n(K)(\hat{g}) = g_*$ which is the chain map induced by the left translation $g: K^H \rightarrow K^H$. This gives a chain complex $\underline{C}_*(K)$ in the abelian category of coefficient systems, and if λ is a coefficient system, then $\text{Hom}(\underline{C}_*(K), \lambda)$, which is the R -module of natural transformations $\underline{C}_*(K) \rightarrow \lambda$, becomes a cochain complex.

Lemma 5. There is an isomorphism of cochain complexes

$$\alpha: C_G^*(K; \lambda) \rightarrow \text{Hom}(\underline{C}_*(K), \lambda).$$

Sketch of Proof. Define α by $\alpha(c)(G/H)(x) = \lambda(G_x \rightarrow H)(c(x))$, where $x \in K_n^H$ and $\lambda(G_x \rightarrow H): \lambda(G/G_x) \rightarrow \lambda(G/H)$ is the homomorphism induced by the inclusion $H \subseteq G_x$. Next, define the inverse α' of α by $\alpha'(T)(x) = T(G/G_x)(x)$. ■

Note that $\underline{C}_*(K)$ is projective in the abelian category of coefficient systems which has sufficiently many injectives, and if λ^* is an injective resolution of λ , then we have a double complex $\text{Hom}(\underline{C}_*(K), \lambda^*)$. The homological algebra applied to this double complex yields a spectral sequence

$$E_2^{p,q} = \text{Ext}^p(\underline{H}_q(K), \lambda) \Rightarrow H_G^{p+q}(K; \lambda),$$

where $\underline{H}_q(K): O_G \rightarrow R\text{-mod}$ is the coefficient system given by

$$\underline{H}_q(K)(G/H) = H_q(K^H; R) \quad \text{and} \quad \underline{H}_q(K)(\hat{g}) = H_q(g).$$

Lemma 6. If $f: K \rightarrow L$ is a G -simplicial map such that each $f^H = f|K^H: K^H \rightarrow L^H$, $H \subseteq G$, induces isomorphism in the classical homology with R coefficients, then

$$f^*: H_G^*(L; \lambda) \rightarrow H_G^*(K; \lambda)$$

is an isomorphism for every coefficient system λ .

Sketch of Proof. We have an isomorphism $f_*: \underline{H}_q(K) \rightarrow \underline{H}_q(L)$ given by $f_*(G/H) = f_*^H$. This extends to an isomorphism f^* between the spectral sequences. ■

For a G -simplicial set K , let RK denote the G -simplicial R -module with the set of n -simplexes $(RK)_n = RK_n$ which is the free R -module with basis K_n and the face and degeneracy maps as the linear extensions of the corresponding maps of K . The G -action on RK is also defined similarly.

Lemma 7. There is an isomorphism $H_G^*(K; \lambda) \cong H_G^*(RK; \lambda)$.

Sketch of Proof. We have a cochain isomorphism $\theta: \text{Hom}(\underline{C}_*(K), \lambda) \rightarrow \text{Hom}(\underline{C}_*(RK), \lambda)$ given by $\theta(T)(G/H)(\sum n_i x_i) = T(G/H)(\sum n_i x_i)$. ■

Let NRK denote the G -pre-simplicial module (degeneracy not considered) where the set of n -simplexes is $\{x \in RK_n : d_i x = 0, 0 \leq i < n\}$, and the n th face operator is d_n .

Lemma 8. There is an isomorphism $H_G^*(K; \lambda) \cong H_G^*(NRK; \lambda)$.

Sketch of Proof. Consider the inclusion map $i: \underline{C}_*(NRK) \rightarrow \underline{C}_*(RK)$. By May [4, (22.3)], $i(G/H): C_*(NRK^H; R) \rightarrow C_*(RK; R)$ induces isomorphism on homology for each $H \subseteq G$. The proof then follows from Lemmas 6 and 7. ■

It may be noted in passing that if X is a G -space and SX the associated singular G -simplicial set, then the cohomology $H_G^*(SX; \lambda)$ is isomorphic to the equivariant singular cohomology of X with coefficient system λ (see Illman [3]), for every λ .

3. Proofs of theorems

Proof of Theorem 1. In view of Lemma 8, it is sufficient to prove that the vector space $C_G^n(NRK; \lambda_G)$ is finitely generated. Let x_1, \dots, x_k denote the representatives of the orbit classes of the non-degenerate n -simplexes which lie in NRK . Suppose that for $1 \leq l \leq k$, the isotropy group G_{x_l} has index m_l in G . Fix a coset representation

$$G/G_{x_l} = \{a_{l_1} G_{x_l}, \dots, a_{l_{m_l}} G_{x_l}\}, \quad 1 \leq l \leq k, \quad a_{l_i} \in G.$$

Then define cochains c_{ij} by

$$c_{ij}(x_l) = \begin{cases} 0 & j \neq l \\ (a_{l_i} G_{x_l})^* & j = l, \quad 1 \leq i \leq m_l \\ 0 & j = l, \quad i > m_l \end{cases}$$

where $(a_{l_i} G_{x_l})^*$ are basis dual to $a_{l_i} G_{x_l}$. There is a unique way to define c_{ij} on the orbit of x_l so that $c_{ij} \in C_G^n(NRK; \lambda_G)$. It is also clear that the set $\{c_{ij}\}$ is a linearly independent set, and that any invariant cochain can be written in terms of the c_{ij} 's. This proves the theorem.

Proof of Theorem 2. The group G is necessarily finite. Therefore $\chi_G(K)$ is defined, and, by Theorem 1, the groups $H_G^*(K; \lambda_{\mathbb{Q}})$ are finitely generated. Also, as the action is free, we have

$$\chi_G(K) = \sum_i (-1)^i N_i = \sum_i (-1)^i \dim_{\mathbb{Q}} H^i(K/G; \mathbb{Q}),$$

where N_i denotes the number of non-degenerate i -simplexes of K modulo the action. Consequently, $\chi_G(K) = \chi(K/G)$, the Euler characteristic of K/G . On the other hand the nature of the action implies $H_G^*(K; \lambda_{\mathbb{Q}}) \cong H^*(K/G; \mathbb{Q}(G))$ and, as

$$\dim_{\mathbb{Q}} H^*(K/G; \mathbb{Q}(G)) = |G| \dim_{\mathbb{Q}} H^*(K/G; \mathbb{Q}),$$

the theorem follows.

Proof of Theorem 3. Since G has finite cohomological dimension, it is torsion free. Also, since G is of finite homological type, the isotropy subgroups G_x also have finite homological type, by a result of Brown [2, IX (6.3)]. Therefore $\chi(G_x)$ is defined, and

$$\begin{aligned} \chi_G(K) &= \sum (-1)^{\dim x} \chi(G_x) = \sum (-1)^{\dim x} \chi(G) [G : G_x] \\ &= \chi(G) \sum_{i=0}^{\dim(K/G)} (-1)^i \dim_{\mathbb{Q}} C_G^i(K; \lambda_{\mathbb{Q}}), \\ &= \chi(G) \sum_{i=0}^{\dim(K/G)} (-1)^i \dim_{\mathbb{Q}} H_G^i(K; \lambda_{\mathbb{Q}}). \end{aligned}$$

The last step follows since we are dealing with vector spaces. This completes the proof.

References

- [1] Bredon G E, *Equivariant cohomology theories*, Springer Lecture Notes in Math. **34** (1967)
- [2] Brown K S, Euler characteristics of discrete groups and G -spaces, *Invent. Math.* **27** (1974) 229–264
- [3] Illman S, Equivariant singular homology and cohomology, *Mem. Am. Math. Soc.* **156** (1975)
- [4] May J P, *Simplicial objects in algebraic topology* (New York: Van Nostrand) (1967)
- [5] Serre J-P, Cohomologie des groupes discrets, *Ann. Math. Stud.* **70** (Princeton: Princeton Univ. Press) (1971)